

Direct extraction of excitation energies from ensemble density-functional theory

Zeng-hui Yang,¹ Aurora Pribram-Jones,^{2,3} Kieron Burke,⁴ and Carsten A. Ullrich⁵

¹*Microsystem and Terahertz Research Center, China Academy of Engineering Physics, Chengdu, China 610200*

²*Department of Chemistry, University of California, Berkeley, CA 94720, USA*

³*Lawrence Livermore National Laboratory, Livermore CA 94550, USA*

⁴*Department of Chemistry, University of California-Irvine, Irvine, CA 92697, USA*

⁵*Department of Physics and Astronomy, University of Missouri, Columbia, MO 65211, USA*

(Dated: April 3, 2017)

A very specific ensemble of ground and excited states is shown to yield an exact formula for any excitation energy as a simple correction to the energy difference between orbitals of the Kohn-Sham ground state. This alternative scheme avoids either the need to calculate many unoccupied levels as in time-dependent density functional theory (TDDFT) or the need for many self-consistent ensemble calculations. The symmetry-eigenstate Hartree-exchange (SEHX) approximation yields results comparable to standard TDDFT for atoms. With this formalism, SEHX yields approximate double-excitations, which are missed by adiabatic TDDFT.

The Hohenberg-Kohn (HK) theorem [1–4] of ground-state density-functional theory (DFT) [1, 5] has several parts. The most-used in practice is the establishment of an exact density functional, $F[n]$, whose minimum yields the exact ground-state density and energy of a given system. Almost all practical calculations use the Kohn-Sham (KS) scheme [5] to minimize F with an approximation to the small exchange-correlation contribution, $E_{xc}[n]$. In fact, many properties of interest in a modern chemical or materials calculation can be extracted from knowledge of the ground-state energy as a function of nuclear coordinates, or in response to a perturbing field.

However, except under very special circumstances, most optical excitation frequencies cannot be deduced. Hence there has always been interest in extending ground-state DFT to include such excitations. Moreover, another part of the HK theorem guarantees that such frequencies (and all properties) are indeed functionals of the ground-state density. In recent years, linear-response time-dependent DFT (TDDFT) [6–10] has become a popular route for extracting low-lying excitation energies of molecules, because of its unprecedented balance of accuracy with computational speed [11]. For significantly sized molecules, more CPU time will be expended on a geometry optimization than a single TDDFT calculation on the optimized geometry.

However, while formally exact, TDDFT with standard approximations is far from perfect. Among the more annoying failures are the underestimation of charge transfer excitations, and the complete lack of multiple excitations when the ubiquitous adiabatic approximation is used [11]. If the unknown exchange-correlation (XC) kernel of TDDFT is approximated by its zero-frequency (and hence ground-state) limit, no multiple excitations survive. While a useful work-around exists for cases where a double is close to a single excitation [12, 13], there is as yet no simple and efficient general procedure for extracting double excitations within adiabatic TDDFT [14].

Ensemble DFT (EDFT) [15, 16] applies the principles

of ground-state DFT to a convex ensemble of the lowest M levels of a system, for which a KS system can be defined [17]. EDFT is formally exact, but practical calculations require approximations, and initial attempts yielded disappointing results [18]. Accuracy is greatly improved when so-called “ghosts” are removed from the approximations [19]. EDFT remains an active research area because, being variational, it should not suffer from some of the limitations of standard TDDFT. Recent strides by Pernal and coworkers [20, 21], Fromager and coworkers [22, 23], and others attempt to create a useful practical alternative to TDDFT, but the difficulty remains in finding accurate low-cost approximations. EDFT usually requires running several different self-consistent ensemble calculations to extract several low-lying excitations.

Here we (a) derive a formula from EDFT to correct a KS orbital energy difference into an exact excitation energy, without doing any self-consistent ensemble calculations, (b) argue that its computational cost should typically be less than either standard TDDFT or EDFT, (c) calculate this correction using the symmetry-eigenstate Hartree-exchange (SEHX) approximation [24, 25], for atoms, demonstrating its accuracy relative to standard TDDFT, and (d) show that SEHX produces estimates of double excitations. This formula, approximation, and the prototype results, may inspire others to explore EDFT more generally.

EDFT is a formally exact and variational excited-state method [15–17]. Let E_i be the electronic energy levels, $i = 0, 1, \dots$, each with degeneracy g_i . Construct an ensemble from positive convex weights w_i , letting I be the maximum non-zero weight. Then, from the foundational theorems, the ensemble energy

$$E_I(\{\mathbf{w}_i\}) = \sum_{i=0}^I g_i w_i E_i, \quad \sum_{i=0}^I g_i w_i = 1, \quad (1)$$

is a functional of the ensemble density

$$n^{\text{ens}}(\mathbf{r}) = \sum_{i=0}^I \mathbf{w}_i n_i(\mathbf{r}), \quad (2)$$

where $n_i(\mathbf{r})$ is the sum of all densities in the i -th multiplet, and can be found via a minimization, so long as the weights are monotonically non-increasing. Applying the same conditions to a fictitious system of non-interacting electrons with the same weights, one can define a KS system whose ensemble density matches the interacting one. Defining energy components in the usual way, only the XC contribution needs to be approximated to perform an ensemble DFT calculation. Since the ensemble energies depend linearly on the weights (at least, in the exact theory), one can easily deduce the transition frequencies.

Infinitely many ensembles can be realized, but the GOK ensemble from the original work [17] is particularly useful and popular, in which all weights are the same except for the highest multiplet, i.e.,

$$\mathbf{w}_{i \neq I} = (1 - g_I \mathbf{w}) / M_{I-1}, \quad \mathbf{w}_I = \mathbf{w}, \quad (\text{GOK}) \quad (3)$$

where M_I is the number of states up to and including the I -multiplet, and $\mathbf{w} \leq M_I^{-1}$ to preserve convexity. When $\mathbf{w} = M_I^{-1}$, the weights are all equal (an equiensemble). In general, the corresponding ensemble density must be found by self-consistent solution of the ensemble KS equations, for the given weights. The excitation energy of the I multiplet can only be isolated by performing self-consistent calculations for all lower multiplets. The excitation energy from the ground state to the I -th multiplet is [17]

$$\omega_I = \frac{1}{g_I} \left. \frac{dE_I^{\text{GOK}}}{d\mathbf{w}} \right|_{\mathbf{w}_I} + \sum_{i=1}^{I-1} \frac{1}{M_i} \left. \frac{dE_i^{\text{GOK}}}{d\mathbf{w}} \right|_{\mathbf{w}_i}, \quad (4)$$

requiring $I + 1$ self-consistent calculations, including the ground-state KS calculation.

The weights defined by Eq. (3) are also a linear interpolation between two consecutive equiensembles, containing M_{I-1} and M_I states. Thus ω_I can also be calculated via

$$\omega_I = M_I E_I(\mathbf{w} = M_I^{-1}) - M_{I-1} E_{I-1}(\mathbf{w} = M_{I-1}^{-1}), \quad (5)$$

which requires only two self-consistent calculations. However, if one needs all excitation energies up to ω_I , $I + 1$ self-consistent calculations are still needed. The computational costs of Eqs. (4) and (5) are much higher than TDDFT with the Casida equation [7].

Now we reintroduce an alternative one-parameter ensemble, in which all states have weight \mathbf{w} *except the ground state*:

$$\mathbf{w}_0 = \frac{1 - \mathbf{w}(M_I - g_0)}{g_0}, \quad \mathbf{w}_{i \neq 0} = \mathbf{w}. \quad (\text{GOKII}) \quad (6)$$

We say reintroduce, as this ensemble was mentioned in a footnote in Ref. [17], although never applied (as far as we know). However, we can show (see supplemental material [26]) that the excitation energy using Eq. (6) has a much simpler formula than using Eq. (4):

$$\omega_I = \frac{1}{g_I} \left[\left. \frac{dE_I^{\text{GOKII}}}{d\mathbf{w}} \right|_{\mathbf{w}_I} - \left. \frac{dE_{I-1}^{\text{GOKII}}}{d\mathbf{w}} \right|_{\mathbf{w}_{I-1}} \right]. \quad (7)$$

Despite the simplicity, in general one still needs to do $I + 1$ calculations to get all excitation energies. However, unlike Eq. (3), the set of weights defined by Eq. (6) is now a linear interpolation between the *ground state* and the equiensemble of M_I states. Now, $\mathbf{w} = 0$ recovers the ground state, not an equiensemble with one less multiplet. A further simplification is made by noting that the EDFT formalism is valid even as $\mathbf{w} \rightarrow 0$. Setting $\mathbf{w}_I = \mathbf{w}_{I-1} = 0$ in Eq. (7) and defining $\Delta\omega_I = \omega_I - \omega_I^{\text{KS}}$, where ω_I^{KS} is the KS orbital energy difference, yields

$$\Delta\omega_I = \frac{1}{g_I} \left. \frac{d}{d\mathbf{w}} \right|_{\mathbf{w}=0} (E_{\text{XC},I}^{\text{GOKII}} - E_{\text{XC},I-1}^{\text{GOKII}}), \quad (\text{DEC}) \quad (8)$$

where $E_{\text{XC},I}^{\text{GOKII}}$ is the GOKII ensemble XC energy functional [17] containing up to the I -th multiplet. This is a direct ensemble correction (DEC) to the KS transition frequencies.

Equation (8) is the central formal result of this work. Because all elements of the right-hand side are evaluated on the ground-state density, this correction is a formally exact ground-state density functional for correcting KS transitions into physical transitions. If approximated by an explicit density functional, it could be evaluated at no noticeable additional cost to a standard ground-state DFT calculation. Compared with the cubic scaling of the TDDFT linear response equations [7], Eq. (8) is vastly more efficient. On the other hand, TDDFT yields both transition frequencies and oscillator strengths, as well as dipole overlap matrices. In addition, linear response TDDFT can yield spatially resolved response functions, once perturbations different from a long-wavelength electric field are allowed. In future work, we will explore what else, beyond transition frequencies, might be extracted in a manner similar to Eq. (8).

There is an infinite number of excited-state ensembles. Even if we consider only those that interpolate between the ground state and the equiensemble, Eq. (6) is not the only choice. The exact ensemble functional yields the same result in *any* ensemble, but approximations yield different results for different ensembles. A DEC expression is a particularly simple route to excitation energies.

Eqs. (3) and (6) are identical for a simple bi-ensemble, the ensemble of a non-degenerate ground and first excited states. Studies of $\mathbf{w} = 0$ bi-ensembles have been carried out previously [27], as well as calculations of the first

excitation energy [24, 25]. Thus the DEC of Eq. (8) can be viewed as a generalization of such results to an arbitrary excitation.

The exact E_{xc}^{GOKII} of Eq. (8) can be obtained numerically for simple cases [24, 25], but in practical calculations E_{xc}^{GOKII} must be approximated. In general, the E_{xc}^{GOKII} of Eq. (8) must account for the state ordering and differences in multiplet structure between the real and KS systems, which poses a challenge for the development of approximations.

SEHX is an explicit orbital-dependent ensemble-density functional analogous to the exact-exchange approximation (EXX) of ground-state DFT. The full expression [24, 25] is long, and given in the supplemental material [26]. In the GOK ensemble, SEHX removes ghost interactions to yield accurate excitation energies [19]. Inserting SEHX into Eq. (8), all the contributions from excitations below I cancel, yielding an approximation that depends only on a difference between a contribution from the I -th multiplet and the ground state:

$$\Delta\omega_I^{\text{SEHX}} = H_I/g_I - H_0/g_0. \quad (9)$$

Here $H_i = H_i^{\text{orb}} + H_i^{\text{dens}}$, where

$$H_i^{\text{orb}} = \frac{1}{2} \int \frac{d^3r d^3r'}{|\mathbf{r} - \mathbf{r}'|} \text{tr} \{ \mathbf{U}(\mathbf{r}, \mathbf{r}') \cdot \mathbf{B}_i + \mathbf{V}(\mathbf{r}, \mathbf{r}') \cdot \mathbf{D}_i \}, \quad (10)$$

and

$$H_i^{\text{dens}} = - \int d^3r v_{\text{HX}}(\mathbf{r}) n_i(\mathbf{r}). \quad (11)$$

Here \mathbf{U} and \mathbf{V} are matrices constructed from KS orbital densities, while \mathbf{B}_i and \mathbf{D}_i are matrices formed from the expression of the spin-symmetric KS eigenfunctions in terms of Slater determinants, as given in the supplemental material, while $v_{\text{HX}}(\mathbf{r})$ is just the ground-state Hartree-exchange potential. We denote calculations with Eq. (9) as DEC/SEHX. Unlike Eq. (8), Eq. (9) depends only on the ground and excited states in question, so the state ordering problem is bypassed and calculation is highly efficient. The ordering-independency of Eq. (9) is due to SEHX, yielding reasonable excitation energies even if the KS state ordering is different from the real one. On the other hand, the approximate state ordering might not be correct.

To illustrate the performance of DEC/SEHX, we calculated excitation energies of small atoms. To see exclusively the effect of the excitation method [28], we use the exact KS potential and energies for the He and Be atoms [29, 30]. We compare with TDDFT using the adiabatic local density approximation (ALDA) [9]. For simplicity, we use the Tamm-Dancoff approximation (TDA) [31] in TDDFT calculations, and we checked to make sure that the results only change slightly with and without TDA. The results are shown in Figs. 1 and 2. More results for atoms are available in the supplemental material [26].




FIG. 1. Exact KS and true excitations of the He atom (black). Experimental values from the NIST atomic spectra database [32, 33]. DEC/SEHX excitation energies in red and TDDFT/ALDA results within TDA in blue.




FIG. 2. Same as Fig. 1, but for Be [32, 34]. Configurations denoted without core. The $2p^2$ configuration corresponds to two doubly-excited states (3^3P and 1^1D).

Figure 1 shows the He results. These are all single excitations (as all doubles in He are in the continuum). The DEC/SEHX gives results that are qualitatively similar to those of standard TDDFT. In fact, the mean absolute errors are typically about 30% *smaller*, despite the lack of approximate correlation in the DEC calculation.

Figure 2 shows the results for Be, again with the exact KS potential. For single excitations, the results are quantitatively similar to those of He, again with DEC er-

rors being noticeably smaller than their TDDFT/ALDA counterparts. But in DEC we can also calculate the double excitations, which are completely absent from any adiabatic TDDFT calculation. We note that the double excitations are less accurate than their single counterparts, but since there are only two, this might be incidental. The supplemental material [26] gives many more atomic calculations, using approximate ground-state KS potentials, showing the strong sensitivity of both DEC and TDDFT to the KS levels in atoms.

To better understand the performance of DEC/SEHX for the double excitations, we turn to a much simpler model problem that was designed to study precisely this question. Consider two fermions in a 1d harmonic potential with a contact interaction [12, 35]:

$$\hat{H} = \frac{1}{2} \sum_{i=1}^2 \left(-\frac{\partial^2}{\partial x_i^2} + x_i^2 \right) + \lambda \delta(x - x'), \quad (12)$$

where $\lambda > 0$. For small values of λ , the system is weakly interacting, and exchange-type approximations are accurate.

		$\Delta\omega_I$				
I	ω_{KS}	exact	TDDFT AEXX	DEC SEHX	exact	TDDFT dressed
Singles						
1	962	38	39	39	38	39
3	1953	47	30	30	48	49
5	2948	52	25	27	51	54
Doubles						
2	1923	41	–	58	41	39
4	2915	49	–	77	49	47

TABLE I. Exact and approximate singlet $\Delta\omega_I$ (in mH) of the 1D two-electron contact-interaction Hooke’s atom with $\lambda = 0.2$. The dressed TDDFT results are calculated with an exchange-only frequency-dependent kernel [12].

The results are shown in Table I. Because this is 1d, there are no degeneracies or multiplets. However, this model was purposely constructed to have near-degeneracies between the multiple and single excitations. With the harmonic confining potential, as $\lambda \rightarrow 0$, many levels approach one another. As shown, the double excitation of level 2 is very close to the single of level 3, and the double at level 4 is very close to the single at level 5.

In the 4th and 5th columns of the table, we report exact exchange results. The former is TDDFT, using the exact KS potential and the exact ground-state exchange in an adiabatic approximation. The latter is DEC/SEHX. We see that both are excellent approximations to the lowest excitations, and give almost identical results for the single excitations. This is because $\lambda = 0.2$, ensuring that correlation effects are relatively weak. But,

unlike adiabatic TDDFT, DEC/SEHX also yields predictions for the double excitations. Just like in the atoms, the errors are substantially larger for the doubles.

Because this model has only two electrons, we can calculate the exact DEC numerically with Eq. (8), by calculating the exact energies, densities and the xc potential of the model first. We then evaluate Eq. (8) numerically using these exact quantities (see supplemental material [26]). These DEC/exact results are in column 6, and agree within a mH with the exact results. This shows that exact DEC does handle doubles correctly, so that the failing in DEC/SEHX is due to the lack of correlation. The last column of the table shows results with the DSPA, a frequency-dependent model XC kernel designed for weakly-correlated systems with strong coupling between a single and double excitation, often called dressed TDDFT [12, 13]. This works extremely well here, as this system was designed to illustrate its accuracy. Study of the difference in the results between these two should provide a route to improving DEC approximations for double excitations.

A discerning reader might have noted that, throughout this work, we have avoided discussion of N - and v -representability[3, 36, 37]. These issues have been partially explored [18, 38] within EDFT in general, but not for this particular ensemble. But none of the calculations here ran into any representation difficulties, such as an inability to find a KS system with the required density. There is little reason to fear such problems in practice. Furthermore, as we use only DEC, any such difficulties in EDFT in general are likely to be least problematic for our applications.

There is obviously much work to be done to see if DEC can become competitive with standard TDDFT calculations. It should be applied to molecules with standard ground-state functionals, to see if the results are as accurate or if semilocal ground-state approximations destroy the accuracy found here within SEHX. Other challenges for TDDFT, such as charge transfer excitations, should be carefully tested. In such a case, we are less hopeful that DEC will provide accurate results as, like TDDFT, it also begins from (unrelaxed) KS transitions of the ground-state. Other ensembles might also yield direct ensemble corrections, or properties other than simple excitation energies might be accessible.

Several other EDFT-based methods for excitations were recently proposed, such as the linear interpolation method [22], the Helmholtz free-energy minimization method [20], and the ensemble-referenced Kohn-Sham method (REKS) [39]. The REKS method is a multi-reference extension to ground-state DFT and EDFT (see also [40]), while the others are within standard EDFT. Each has its own advantages, and the REKS method has been shown to work well in strongly-correlated systems [41]. However, all these methods require extra self-consistent calculations aside from the ground-state one.

The simplification achieved in this paper by changing the ensemble type suggests that similar simplifications may also be possible in these methods.

But for now, DEC, as presented here in Eq. (8), is a promising alternative to existing approaches to excitation energies from DFT. Relative to standard TDDFT, it has lower computational cost and is applicable to traditionally difficult problems such as multiple excitations and spin-multiplets. Unlike TDDFT, EDFT is based on a variational principle[16], so the DEC derived in this work may be more reliable than TDDFT corrections, which are based on response theory. With the DEC, the calculation of excitation energies can be carried out efficiently as a post-processing step after a ground-state KS calculation. The calculations shown in this paper are merely intended to demonstrate the DEC method. The SEHX approximation, while showing better accuracy than TDDFT/ALDA for single excitations in atoms and producing double excitations, does not yield as high accuracy for the doubles. We stress that such limitations are due to the ordering-independent SEHX approximation only. The DEC derived in this work is formally exact, as illustrated in our model harmonic trap.

The authors thank Cyrus Umrigar for providing us the exact KS potentials of the He and Be atoms. Z.-H.Y. and C.A.U. were supported by NSF grant DMR-1408904. A.P.J. was supported by the University of California President's Postdoctoral Fellowship. Part of this work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344. K.B. was supported by DOE grant DE-FG02-08ER46496. Z.-H.Y. is currently supported by the Science Challenge Project (China) grant No. JCKY2016212A503-1.

-
- [1] P. Hohenberg and W. Kohn, Phys. Rev. **136**, B864 (1964).
- [2] M. Levy, Proc. Natl. Acad. Sci. USA **76**, 6062 (1979).
- [3] M. Levy, Phys. Rev. A **26**, 1200 (1982).
- [4] E. H. Lieb, Int. J. Quantum Chem. **24**, 243 (1983).
- [5] W. Kohn and L. J. Sham, Phys. Rev. **140**, A1133 (1965).
- [6] E. Runge and E. K. U. Gross, Phys. Rev. Lett. **52**, 997 (1984).
- [7] M. E. Casida, in *Recent Developments and Applications of Modern Density Functional Theory*, Theoretical and Computational Chemistry, Vol. 4, edited by J. M. Seminario (Elsevier, Amsterdam, 1996) p. 391.
- [8] M. A. L. Marques, N. T. Maitra, F. M. S. Nogueira, E. K. U. Gross, and A. Rubio, eds., *Fundamentals of Time-Dependent Density Functional Theory*, Lecture Notes in Physics (Springer, Berlin, 2012).
- [9] C. A. Ullrich, *Time-Dependent Density-Functional Theory: Concepts and Applications* (Oxford University Press, Oxford, 2012).
- [10] C. A. Ullrich and Z.-H. Yang, Brazilian J. Phys. **44**, 154 (2014).
- [11] N. Maitra, J. Chem. Phys. **144**, 220901 (2016).
- [12] N. T. Maitra, F. Zhang, R. J. Cave, and K. Burke, J. Chem. Phys. **120**, 5932 (2004).
- [13] R. J. Cave, F. Zhang, N. T. Maitra, and K. Burke, Chem. Phys. Lett. **389**, 39 (2004).
- [14] P. Elliott, S. Goldson, C. Canahui, and N. T. Maitra, Chem. Phys. **391**, 110 (2011).
- [15] A. Theophilou, J. Phys. C **12**, 5419 (1979).
- [16] E. K. U. Gross, L. N. Oliveira, and W. Kohn, Phys. Rev. A **37**, 2805 (1988).
- [17] E. K. U. Gross, L. N. Oliveira, and W. Kohn, Phys. Rev. A **37**, 2809 (1988).
- [18] L. N. Oliveira, E. K. U. Gross, and W. Kohn, Phys. Rev. A **37**, 2821 (1988).
- [19] N. I. Gidopoulos, P. G. Papaconstantinou, and E. K. U. Gross, Phys. Rev. Lett. **88**, 033003 (2002).
- [20] E. Pastorczak, N. I. Gidopoulos, and K. Pernal, Phys. Rev. A **87**, 062501 (2013).
- [21] E. Pastorczak and K. Pernal, J. Chem. Phys. **140**, 18A514 (2014).
- [22] B. Senjean, S. Knecht, H. J. A. Jensen, and E. Fromager, Phys. Rev. A **92**, 012518 (2015).
- [23] K. Deur, L. Mazouin, and E. Fromager, Phys. Rev. B **95**, 035120 (2017).
- [24] A. Pribram-Jones, Z.-H. Yang, J. R. Trail, K. Burke, R. J. Needs, and C. A. Ullrich, J. Chem. Phys. **140**, 18A541 (2014).
- [25] Z.-H. Yang, J. R. Trail, A. Pribram-Jones, K. Burke, R. J. Needs, and C. A. Ullrich, Phys. Rev. A **90**, 042501 (2014).
- [26] See supplemental material at [url].
- [27] M. Levy, Phys. Rev. A **52**, R4313 (1995).
- [28] M.-C. Kim, E. Sim, and K. Burke, Phys. Rev. Lett. **111**, 073003 (2013).
- [29] C. J. Umrigar and X. Gonze, in *High Performance Computing and its Applications in the Physical Sciences*, Proceedings of the Mardi Gras '93 Conference, edited by D. A. Browne (World Scientific, Singapore, 1994).
- [30] C. J. Umrigar and X. Gonze, Phys. Rev. A **50**, 3827 (1994).
- [31] S. Hirata and M. Head-Gordon, Chem. Phys. Lett. **314**, 291 (1999).
- [32] <http://www.nist.gov/pml/data/asd.cfm>.
- [33] D. C. Morton, Q. Wu, and G. W. F. Drake, Can. J. Phys. **84**, 83 (2006).
- [34] A. Kramida and W. C. Martin, J. Phys. Chem. Ref. Data **26**, 1185 (1997).
- [35] F. Zhang and K. Burke, Phys. Rev. A **69**, 052510 (2004).
- [36] R. G. Parr and W. Yang, *Density Functional Theory of Atoms and Molecules* (Oxford University Press, 1989).
- [37] T. L. Gilbert, Phys. Rev. B **12**, 2111 (1975).
- [38] N. Hadjisavvas and A. Theophilou, Phys. Rev. A **32**, 720 (1985).
- [39] M. Filatov, WIREs Comput. Mol. Sci. **5**, 146 (2015).
- [40] E. Pastorczak and K. Pernal, Int. J. Quant. Chem. **116**, 880 (2016).
- [41] M. Filatov, M. Huix-Rotllant, and I. Burghardt, J. Chem. Phys. **142**, 184104 (2015).