

Reassessing the role of exact conditions in density functional theory

Ryan Pederson*

Department of Physics and Astronomy, University of California, Irvine, CA 92697, USA

Kieron Burke†

Department of Chemistry, University of California, Irvine, CA 92697, USA and

Department of Physics and Astronomy, University of California, Irvine, CA 92697, USA

(Dated: March 6, 2023)

Exact conditions have long been used to guide the construction of density functional approximations. Nowadays hundreds of approximations are in common use, many of which neglect these conditions in their design. We analyze several well-known exact conditions and revive several obscure ones. Two crucial distinctions are drawn: that between necessary and sufficient conditions, and between all possible electronic densities and the subset of relevant Coulombic ground states. Simple search algorithms find violations of sufficient conditions while others construct densities that violate necessary conditions. We find that many empirical approximations satisfy many exact conditions for chemically relevant densities. We also find non-empirical approximations satisfy even more conditions than those enforced in their construction.

Modern density functional theory (DFT) calculations span many branches of the science of matter [1–4]. In the standard Kohn-Sham approach [5], only the exchange-correlation (XC) energy need be approximated as a functional of the electronic (spin)-densities. Currently, hundreds of distinct XC approximations are available in standard DFT codes [6, 7], reflecting the immense difficulty in finding approximations that are generally accurate.

With the exact XC energy functional, Kohn-Sham DFT provides the exact ground-state density and energy for any electronic system. Exact conditions are known analytical properties of the exact functional and have played a vital role in the development of approximations [8]. The argument is that imposing exact conditions makes an approximation better resemble the exact functional, leading to improved generality. Typically, *non-empirical* functionals rely heavily on such conditions, eschew fitting to any chemical bonds, and work reasonably well for both materials and molecules. Such guiding principles led to a series of successful and widely used approximations, culminating in “Strongly Constrained and Appropriately Normed” (SCAN) [9] semilocal functional which attributes much of its success to the satisfaction of ‘all known’ (17) exact conditions that such a functional can satisfy.

On the other hand, many approximations tailored for molecular chemistry applications blatantly ignore exact conditions in their design [10]. Such approximations can be extremely accurate on comprehensive molecular benchmarks [11], where they are often more accurate than their more constrained counterparts. Typically, such chemically trained functionals behave poorly for materials.

We illustrate this difference with the correlation energy of the blue two-electron density in Fig 1, calculated with two generalized gradient approximations (GGA’s). The first, the Perdew-Burke-Ernzerhof (PBE) correlation functional [12]




FIG. 1. An unpolarized ground-state gedanken density with 2 electrons whose correlation energy is -21 mH in PBE, but +85 mH in LYP. For reference, the He atom density (divided by 7) is plotted.

adheres to many exact conditions and automatically satisfies the basic requirement that the correlation energy is never positive, yielding -21 mH. The second, the Lee-Yang-Parr (LYP) correlation functional [13], does not explicitly enforce many exact conditions and yields the nonsensical +85 mH. Yet LYP has been used successfully in over 100,000 chemical applications [14].

How can the success of these two seemingly disjoint design paradigms be rationalized? We resolve this paradox by reassessing the role of exact conditions in modern DFT approximations. To do this, we develop several new (and not so new) tools. We carefully parse the logic of exact conditions, finding that many enforced conditions are too strong for real matter. A computational scanning procedure finds violations, coupled with construction of corresponding reasonable (but not realistic) densities, as in Fig 1. Half

* pedersor@uci.edu

† kieron@uci.edu

a dozen exact conditions and hundreds of approximate functionals are analyzed. Several obscure conditions are revived and analyzed, while even well-known conditions yield surprising new twists. Finally, the role of exact conditions in density functional development is revisited.

Begin with the correlation energy. In practice, approximations (denoted by tilde) have the form:

$$\tilde{E}_C[n] = \int d^3r n(\mathbf{r}) \tilde{\epsilon}_C[n](\mathbf{r}), \quad (1)$$

where $n(\mathbf{r})$ is an electronic density. While the developers define a conventional correlation energy per electron, $\tilde{\epsilon}_C[n](\mathbf{r})$, that is often implemented explicitly in DFT codes, other “gauges” exist yielding the same $\tilde{E}_C[n]$. For example, $\tilde{\epsilon}_C[n](\mathbf{r})$ and $\tilde{\epsilon}_C[n](\mathbf{r}) + \nabla^2(n^{2/3})/n$ yield identical $\tilde{E}_C[n]$ [15, 16]. Specifically, semilocal functionals can be written

$$\tilde{\epsilon}_C[n](\mathbf{r}) = \tilde{\epsilon}_C(r_s(\mathbf{r}), \zeta(\mathbf{r}), s(\mathbf{r}), \alpha(\mathbf{r}), q(\mathbf{r})), \quad (2)$$

where $r_s = (4\pi n/3)^{-1/3}$ is the Wigner-Seitz radius, $\zeta = (n_\uparrow - n_\downarrow)/n$ is the (dimensionless) spin polarization, $s = |\nabla n|/(2(3\pi^2)^{1/3}n^{4/3})$ is the (dimensionless) reduced gradient, $\alpha = (\tau - \tau^{VW})/\tau^{\text{unif}} \geq 0$ with $\tau = \sum_{i,\sigma}^{\text{occ.}} |\nabla \phi_{i,\sigma}|^2/2$, $\tau^{VW} = |\nabla n|^2/8n$, $\tau^{\text{unif}} = (3/20)(3\pi^2)^{2/3}n^{5/3}[(1 + \zeta)^{5/3} + (1 - \zeta)^{5/3}]$, and $q = \nabla^2 n/(4(3\pi^2)^{2/3}n^{5/3})$ is the reduced Laplacian. The local spin density approximations (LDA) depends only on r_s and ζ , generalized gradient approximations (GGAs) add dependence on s , while meta-generalized gradient approximations (MGGAs) can depend on all variables.

A simple exact condition is *correlation energy non-positivity*,

$$E_C[n] \leq 0, \quad (3)$$

which holds for any reasonable density, which we define as being positive, integrating to a finite quantity N , and have finite von Weizsäcker kinetic energy (\mathcal{I}_N of Ref. [17] or Eq. 34 of Ref. [18]). This is routinely enforced via

$$\tilde{\epsilon}_C[n](\mathbf{r}) \leq 0 \quad \text{for all } \mathbf{r} \text{ and any } n(\mathbf{r}). \quad (4)$$

Clearly, satisfying this local condition *guarantees* Eq. (3), but it is also excessive, i.e., not necessary. Moreover, starting from any $\tilde{\epsilon}_C[n](\mathbf{r})$ that satisfies Eq (4), addition of $C\nabla^2(n^{2/3})/n$ violates it for sufficiently large C . If local violations of Eq. (4) *do* exist, then a counterexample density that violates the exact condition in Eq. (3) *might* be found. If it can be, the exact condition is violated for that density in any gauge. If no such counterexample can be found, the possibility that a gauge might be found that satisfies Eq. (4) remains open.

Returning to the LYP GGA, we found instances where $\epsilon_C^{\text{LYP}}(r_s, \zeta, s) > 0$ for $s \geq 1.74$, thus allowing the possibility of a violating *gedanken* density. Gedanken densities are thought experiment densities that need not be realistic [15]. So we construct a *gedanken* density that has large $s \geq 1.74$

values throughout its interior, violating the local condition in Eq. (4) (see Supplemental S1 for details). Importantly, we want such local violations to exist in *energetically relevant regions* of the density, that is, spatial regions that substantially contribute to the integral in Eq. 1. The *gedanken* density of Fig 1 is radial, nodeless, finite, continuous, differentiable to first and second order, and integrates to 2 electrons. It is a reasonable density and is also non-interacting v -representable, and when evaluated using the LYP correlation functional yields +85 mH. Thus, the LYP functional *can* violate correlation non-positivity.

But does LYP violate correlation non-positivity in the restricted space of realistic Coulombic densities, i.e, those ground-state densities of systems with Coulombic attractions to integer nuclear charges of small or no overall charge? The *gedanken* density of Fig 1 is not Coulombic: for instance, it lacks nuclear cusps as required by Kato’s theorem [19].




FIG. 2. The distribution $g(s)$ is plotted for various ground-state densities: the *gedanken* density in Fig. 1, the He and N atoms, and the N_2 molecule. The absolute difference between the N_2 molecule and N atom distributions is also plotted. Details of the calculations can be found in Supplemental S2.

An important property of the *gedanken* density is that it has energetically relevant regions of the density with $s \geq 2$. The distribution

$$g(s) = \int d^3r n(\mathbf{r}) \delta^{(3)}(s - s(\mathbf{r})), \quad (5)$$

was introduced in Refs. [20, 21], and $g(s) ds$ is the number of electrons in the system with reduced density gradient between s and $s + ds$, i.e, it is an analog of the density of states for energy levels. In Fig. 2, we plot $g(s)/N$ for various ground-state densities: the *gedanken* density in Fig. 1 and calculated densities for the He and N atoms and the N_2 molecule. Hartree-Fock densities are sufficiently accurate for our purposes. Unlike the Coulombic densities, the *gedanken* density $g(s)$ is centered around $s \approx 2$, as

intended. For Coulombic systems [21], large $s > 2$ values are typically only found in the decaying tails of the density, and are energetically irrelevant. In molecular and extended systems, these tails (which may not even be present in periodic systems) are even less energetically relevant than their atomized counterparts [21]: in Fig. 2 the distribution $g(s)/N$ of N_2 is shifted to lower s than that of the atomized system (N atom) for $s > 1$. The electrons in the bond between two (or more) atoms have smaller values of s which even vanishes at the bond center [21]. Although a single electron distributed across an infinitely separated chain of many protons has energetically relevant regions of large s , such a system is too far from neutral for our set.

Do large s values contribute importantly to energy differences, such as molecular binding energies? Valence electrons change considerably but their differences tend to also be energetically relevant only in regions of smaller s values: the difference $|g[N_2](s) - 2g[N](s)|$, is most prominent in regions $s < 2$ (see Fig. 2) [21]. In non-covalent bonding, s values up to ≈ 7 are relevant in binding energy differences for van der Waals (vdW) complexes [22, 23]. However, in these cases, typically a non-local correlation functional (such as DFT-D [24] or vdW-DF [25]) is employed which provides the bulk of the energetics that contributes to the binding of vdW complexes, whereas semilocal contributions are more relevant in structural aspects such as bonding distances and lattice constants [22].

Revisiting the LYP correlation energy example, we conjecture that no realistic Coulombic density ever yields a positive correlation energy. Such a density would need to have energetically relevant regions of the density with large $s > 1.74$, which is simply not observed in these systems. Over countless atomic and molecular densities, the LYP correlation functional has *not* yielded positive correlation energies [13, 26].

We perform an identical analysis on other popular approximations and tabulate the results in the first row of Table I. For each exact condition (e.g. Eq. (3)), we check for violations of the corresponding local condition (e.g. Eq. (4)) for various semilocal approximations. To locate violations of such local conditions, we use a simple extensive grid search over applicable variables, $0 < r_s \leq 5, 0 \leq \zeta \leq 1, 0 \leq s \leq 5, 0 \leq \alpha \leq 5$, where the upper ranges are chosen to be reasonably large to encapsulate relevant regions of realistic Coulombic densities (see Supplemental S11 for more details). This brute-force numerical approach is necessary since analytical verification is often impossible with the complicated forms of modern XC approximations. We utilize the Libxc [7] implementation of XC approximations to evaluate the XC energy densities in Eq. (2).

For a given local condition and approximation, if we find no violations in our grid search, then we assume that the approximation always satisfies the exact condition for any reasonable density (denoted by a \checkmark in Table I). In other cases (denoted \checkmark^*), the corresponding local condition may be satisfied only for a restricted range of variable values. For example, for B3LYP correlation non-positivity, the local

condition in Eq. 4 is satisfied whenever $s < 2.13$ and we display the bounds on the variable s that ensure satisfaction. In some cases (denoted \checkmark^{**}), we do not obtain a simple restricted range of variable values, but we find that local violations are exceedingly “rare” (less than 1% of the total number of permutations considered).

The logic and concepts presented for correlation non-positivity generalize to other exact conditions on energy functionals. For instance, the *Lieb-Oxford (LO) bound* [27] *extension to XC energies* exact condition [28],

$$E_{\text{XC}}[n] \geq C_{\text{LO}} \int d^3r n \epsilon_{\text{x}}^{\text{unif}}[n](\mathbf{r}), \quad (6)$$

yields the local condition

$$\tilde{F}_{\text{XC}} \leq C_{\text{LO}}, \quad (7)$$

where we use $C_{\text{LO}} = 2.27$ in this work (although tighter bounds have been proven [29, 30]), $\tilde{F}_{\text{XC}}[n](\mathbf{r}) \equiv \tilde{\epsilon}_{\text{XC}}[n](\mathbf{r})/\epsilon_{\text{x}}^{\text{unif}}[n](\mathbf{r})$ is the exchange (correlation) enhancement factor with $\tilde{F}_{\text{XC}} = \tilde{F}_{\text{x}} + \tilde{F}_{\text{c}}$ and $\epsilon_{\text{x}}^{\text{unif}}[n](\mathbf{r}) = -(3/4\pi)(3\pi^2n)^{1/3}$ is the exchange energy per particle of an unpolarized uniform electron gas. Many approximations enforce the local Eq. (7) to ensure Eq. 6.

Since the combined XC energy is the object of interest, some approximations fail to distinguish exchange and correlation. In the exact functional, one can extract E_{C} using uniform coordinate scaling [31, 32]:

$$E_{\text{C}}[n] = E_{\text{XC}}[n] - \lim_{\gamma \rightarrow \infty} \frac{E_{\text{XC}}[n_{\gamma}]}{\gamma}, \quad (8)$$

where $n_{\gamma}(\mathbf{r}) \equiv \gamma^3 n(\gamma \mathbf{r})$ and $\gamma > 0$. We apply this “conventional” partitioning to extract correlation energies where none have been defined or whose partitioning is ambiguous. E.g., for globals, Eq. (8) yields E_{C} of the semilocal form in Eq. (2), e.g., $\epsilon_{\text{C}}^{\text{B3LYP}} = 0.405\epsilon_{\text{C}}^{\text{LYP}} + 0.095\epsilon_{\text{C}}^{\text{VWN5}}$ [33]. This partitioning can differ from the developer’s intentions or rationalizations (see Supplemental S4). The LO bound applied to global hybrid functionals is discussed in Supplemental S5.

Besides Eq. (8), many other properties of the exact functional are written in terms of uniform coordinate scaling (or equivalently, through the adiabatic connection in DFT [34–36], see Supplemental S7). We simply list these “obscure” conditions and their local forms (further details and derivations can be found in the Supplemental S3). The *correlation uniform scaling inequality* [31]

$$(\gamma - 1)E_{\text{C}}[n_{\gamma}] \geq \gamma(\gamma - 1)E_{\text{C}}[n] \quad (9)$$

has a corresponding local condition,

$$\frac{\partial \tilde{F}_{\text{C}}(r_s, \zeta, s, \alpha, q)}{\partial r_s} \geq 0. \quad (10)$$

The kinetic contribution to the correlation energy, T_{C} , is non-negative [31, 37]

$$T_{\text{C}}[n_{\gamma}] = \gamma \frac{dE_{\text{C}}[n_{\gamma}]}{d\gamma} - E_{\text{C}}[n_{\gamma}] \geq 0, \quad (11)$$

local condition	non-empirical			empirical				
	PBE	AM05	SCAN	B3LYP	CASE21	SOGGA11	M06	B97
E_C non-positivity (4)	✓	✓	✓	✓*, $s < 2.13$	✓	✓	✓*, $\zeta = 0, s < 1.56$	✓*, $s < 1.42$
E_C scaling inequality (10)	✓	✓	✓	✓*, $s < 2.15$	✓	✓**	✓*, $\zeta = 0, s < 1.59$	✓*, $s < 1.52$
T_C upper bound (13)	✓	✓	✓	✓	✓	✓*, $s < 1.36$	✓*, $\zeta = 0, s < 1.56$	✓*, $s < 1.62$
$U_C(\lambda)$ monotonicity (15)	✓	✓	✓	✓*, $s < 1.82$	✓	✗	✓*, $\zeta = 0, s < 1.56$	✓*, $s < 1.41$
LO extension to E_{XC} (7)	✓	✓	✓	✓	✓	✓	✓*, $0.04 < s < 3.62$	✓*, $s < 4.46$
LO (17)	✓	✓	✓	✓*, $s < 4.88$	✓	✓*, $s < 4.98$	✓*, $0.06 < s < 3.62$	✓*, $s < 4.43$

TABLE I. For each condition, we assess if the local condition is satisfied (or partially satisfied) for an approximation (with more given in Supplemental S11).

and shares the same local condition, Eq. 10. The T_C upper bound [38, 39] reads

$$T_C[n_\gamma] \leq -\gamma \left(\frac{\partial E_C[n_\gamma]}{\partial \gamma} \Big|_{\gamma \rightarrow 0} \right) + E_C[n_\gamma], \quad (12)$$

with corresponding local condition,

$$\frac{\partial \tilde{F}_C}{\partial r_s} \leq \frac{\tilde{F}_C(\infty) - \tilde{F}_C}{r_s}, \quad (13)$$

where $\tilde{F}_C(\infty) = \tilde{F}_C(r_s \rightarrow \infty)$. Correlation energy adiabatic connection curves, $U_C(\lambda) = d(\lambda^2 E_C[n_{1/\lambda}])/d\lambda$, satisfy a *monotonicity condition* [32],

$$\frac{dU_C(\lambda)}{d\lambda} \leq 0, \quad (14)$$

with corresponding local condition [32]

$$\frac{\partial}{\partial r_s} \left(r_s^2 \frac{\partial \tilde{F}_C}{\partial r_s} \right) \geq 0. \quad (15)$$

The LO bound [27], often generalized as Eq. 6, is precisely

$$U_{XC}[n] \geq C_{LO} \int d^3r n \epsilon_x^{\text{unif}}[n](\mathbf{r}), \quad (16)$$

where $U_{XC}[n] = E_{XC}[n] - T_C[n]$ is the potential correlation energy. The corresponding local condition,

$$\tilde{F}_{XC} + r_s \frac{\partial \tilde{F}_C}{\partial r_s} \leq C_{LO}, \quad (17)$$

is *more* restrictive than the commonly used Eq. (7). Approximations satisfying Eq. (7) need not satisfy Eq. (17), such as B3LYP or SOGGA11 in Table I. Results for a conjectured condition, $T_C[n] \leq -E_C[n]$ [31, 37, 40], are in Supplemental S6, S10, and S11.

This work does not provide a comprehensive study of all known exact conditions in DFT. A unified subset

of several conditions (6) were chosen to illustrate the logic. But we have not touched on the self-interaction error [41], the asymptotic behavior of exchange and correlation potentials [42], or the flat-plane energy condition for fractional charges and spins [43, 44]. We expect our logic can be fruitfully applied to any exact condition in DFT.

We describe the conditions Eqs. 9 - 17 as obscure because, while proven several decades ago, none appear to be deliberately and generally enforced in modern approximations, even those that strive to satisfy as many exact conditions as possible. SCAN was designed to satisfy the correlation uniform scaling inequality (Eq. (9)), but only in extreme limits, $\gamma \rightarrow 0, \infty$ [8, 9]. The corresponding local condition in Eq. (10) is satisfied in SCAN, but adjustments of its parameters chosen to fit *appropriate norms* can produce violations (see Supplemental S9). Norms refer to properties of specific (but not bonded) reference systems, such as the uniform gas, the hydrogen atom, or noble gas dimers [45, 46]. So, by enforcing appropriate norms, SCAN satisfies *more* exact conditions than were explicitly included. In other instances, such as PW91 [28] and LDA, the satisfaction of several of the obscure exact conditions was proven analytically after their publication [32] (Supplemental S8 is a proof that PBE satisfies Eq. 9). We find (numerically) that the non-empirical PBE, AM05, and SCAN functionals, which were designed to satisfy a large set of exact conditions, also satisfy many additional exact conditions outside of the original set (Table I). CASE21, a recent machine-learned empirical functional designed to adhere to select exact conditions [47], also satisfies these obscure conditions.

In Table I, we also show that many empirical approximations satisfy local conditions in energetically relevant regions of realistic Coulombic densities, i.e., for s values that are not too large. When assessing our set of exact conditions on the set of coordinate-scaled HF densities for atoms H-Ar and their cations, we find that *all* are satisfied (see Supplemental S10), suggesting that these approximations will satisfy these conditions

in the space of realistic Coulombic densities (possibly excepting the monotonicity condition in SOGGA11). This is intriguing because most such empirical approximations were designed *without* explicit adherence to these exact conditions. This finding appears to reinforce their importance in approximations: satisfaction of such esoteric exact conditions is hardly accidental. Furthermore, empirical approximations often employ ingredients, such as the dimensionless quantities s, ζ, α, q , which themselves were chosen to simplify satisfaction of exact conditions. In consequence, nearly all empirical approximations satisfy two simple exact constraints on the exchange energy: uniform coordinate scaling [31] and spin scaling [48].

Our results suggest a reassessment of the role of exact conditions in modern density functional development. Is it excessive to enforce strong local conditions to ensure the satisfaction of exact conditions? Flexible empirical model approximations also satisfy many exact conditions on the (highly relevant) space of realistic Coulombic densities and achieve high accuracy for molecular processes. When empirical functionals are locally constrained to satisfy conditions, their molecular benchmark performance is similar to the suboptimal performance of their non-empirical counterparts, despite the advantage of training on molecular data [47, 49, 50], and the resulting functionals tend to closely mimic non-empirical counterparts, such as SCAN [49, 50].

So when is the enforcement of local conditions helpful? The answer appears to lie in the paucity of highly accurate data for solids. Empirical approximations are mostly developed for molecules, where copious benchmark data is now available, and are rarely even available in materials codes (as they typically perform poorly for solids). On the other hand, non-empirical functionals that dominate materials calculations include norms such as the uniform gas limit, which take the place of highly accurate data. By reducing to this limit, non-empirical functionals are guaranteed to yield moderately accurate results for solids. Moreover, the leading corrections in the asymptotic limit differ qualitatively between molecules and solids, because all molecules have turning surfaces at the Kohn-Sham HOMO energy, while few solids do [51]. This produces conflicting requirements on the gradient expansion of the approximation, as shown in the differences between PBE (good for atomic and molecular energies) and PBEsol (good for solid geometries and vibrations [52]). Such conflicts are resolved in SCAN, yielding improved results for both. Even in the presence of highly accurate data for solids, approximations that fit their data will worsen their results for molecules.

Even for molecular systems, non-empirical approximations like SCAN may also outperform empirical approximations with the same ingredients, especially when applied to new systems or new properties outside their training [8, 53, 54]. For instance, when evaluated on the artificial molecules in the MB16 benchmark [11, 55] (never used to parameterize empirical functionals), SCAN tends to outperform empirical approximations, including hybrids [8],

suggesting a greater ability to extrapolate.

In conclusion, this work explores and revives several exact conditions in DFT and provides analysis on their satisfaction in approximations. Empirical-based approximations, which often employ flexible models that do not explicitly enforce exact conditions, typically violate exact conditions on some densities, but do not violate them for realistic Coulombic systems. As such approximations are trained on such systems, relevant exact conditions are implicitly enforced. Specific exact conditions (and appropriate norms) can be especially important when lacking reference data, where their enforcement (either approximate or exact) helps maintain the transferability of resulting approximations, e.g., from molecules to solids. These considerations may aid future approximations to achieve high accuracy that is maintained across many different classes of systems: molecules, solids, and everything in between, such as interfaces, clusters, and metal-organic frameworks.

1. ACKNOWLEDGMENTS

Work supported by DOE DE-SC0008696. We thank John P. Perdew and Aaron D. Kaplan for useful discussions.

- [1] J. K. Nørskov, F. Abild-Pedersen, F. Studt, and T. Bligaard, *Proceedings of the National Academy of Sciences* **108**, 937 (2011), <https://www.pnas.org/doi/pdf/10.1073/pnas.1006652108>.
- [2] A. Jain, Y. Shin, and K. A. Persson, *Nature Reviews Materials* **1**, 1 (2016).
- [3] C. J. Pickard, I. Errea, and M. I. Eremets, *Annual Review of Condensed Matter Physics* **11**, 57 (2020).
- [4] L. Zeng, S. B. Jacobsen, D. D. Sasselov, M. I. Petaev, A. Vanderburg, M. Lopez-Morales, J. Perez-Mercader, T. R. Mattsson, G. Li, M. Z. Heising, *et al.*, *Proceedings of the National Academy of Sciences* **116**, 9723 (2019).
- [5] W. Kohn and L. J. Sham, *Phys. Rev.* **140**, A1133 (1965).
- [6] N. Mardirossian and M. Head-Gordon, *Molecular Physics* **115**, 2315 (2017).
- [7] S. Lehtola, C. Steigemann, M. J. Oliveira, and M. A. Marques, *SoftwareX* **7**, 1 (2018).
- [8] A. D. Kaplan, M. Levy, and J. P. Perdew, *arXiv preprint arXiv:2207.03855* (2022).
- [9] J. Sun, A. Ruzsinszky, and J. P. Perdew, *Phys. Rev. Lett.* **115**, 036402 (2015).
- [10] R. Peverati and D. G. Truhlar, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* **372**, 20120476 (2014).
- [11] L. Goerigk, A. Hansen, C. Bauer, S. Ehrlich, A. Najibi, and S. Grimme, *Physical Chemistry Chemical Physics* **19**, 32184 (2017).
- [12] J. P. Perdew, K. Burke, and M. Ernzerhof, *Phys. Rev. Lett.* **77**, 3865 (1996), *ibid.* **78**, 1396(E) (1997).
- [13] C. Lee, W. Yang, and R. G. Parr, *Phys. Rev. B* **37**, 785 (1988).
- [14] K. Burke, *J. Chem. Phys.* **136** (2012).
- [15] J. P. Perdew, A. Ruzsinszky, J. Sun, and K. Burke, *The Journal of Chemical Physics* **140**, 18A533 (2014).
- [16] F. G. Cruz, K.-C. Lam, and K. Burke, *The Journal of Physical Chemistry A* **102**, 4911 (1998).
- [17] E. H. Lieb, *Int. J. Quantum Chem.* **24**, 243 (1983).
- [18] L. O. Wagner, T. E. Baker, M. Stoudenmire, E., K. Burke, and S. R. White, *Phys. Rev. B* **90**, 045109 (2014).
- [19] T. Kato, *Communications on Pure and Applied Mathematics* **10**, 151 (1957).
- [20] A. Zupan, J. P. Perdew, K. Burke, and M. Causà, *International journal of quantum chemistry* **61**, 835 (1997).
- [21] A. Zupan, K. Burke, M. Ernzerhof, and J. P. Perdew, *The Journal of chemical physics* **106**, 10184 (1997).
- [22] T. Jenkins, K. Berland, and T. Thonhauser, *Electronic Structure* **3**, 034009 (2021).
- [23] É. D. Murray, K. Lee, and D. C. Langreth, *Journal of Chemical Theory and Computation* **5**, 2754 (2009).
- [24] D. Chakraborty, K. Berland, and T. Thonhauser, *Journal of Chemical Theory and Computation* **16**, 5893 (2020).
- [25] S. Grimme, J. Antony, S. Ehrlich, and H. Krieg, *The Journal of chemical physics* **132**, 154104 (2010).
- [26] B. Miehlisch, A. Savin, H. Stoll, and H. Preuss, *Chemical Physics Letters* **157**, 200 (1989).
- [27] E. H. Lieb and S. Oxford, *International Journal of Quantum Chemistry* **19**, 427 (1981).
- [28] J. P. Perdew, in *Proceedings of the 75. WE-Heraeus-Seminar and 21st Annual International Symposium on Electronic Structure of Solids*, edited by P. Ziesche and H. Eschrig (Akademie Verlag, Berlin, 1991) p. 11.
- [29] M. Lewin, E. H. Lieb, and R. Seiringer, *Phys. Rev. B* **100**, 035127 (2019).
- [30] M. Lewin, E. H. Lieb, and R. Seiringer, *Improved lieb-oxford bound on the indirect and exchange energies* (2022).
- [31] M. Levy and J. P. Perdew, *Physical Review A* **32**, 2010 (1985).
- [32] M. Levy and J. P. Perdew, *Physical Review B* **48**, 11638 (1993).
- [33] P. J. Stephens, F. J. Devlin, C. F. Chabalowski, and M. J. Frisch, *The Journal of physical chemistry* **98**, 11623 (1994).
- [34] J. Harris and R. Jones, *J. Phys. F* **4**, 1170 (1974).
- [35] D. Langreth and J. Perdew, *Solid State Commun.* **17**, 1425 (1975).
- [36] O. Gunnarsson and B. Lundqvist, *Phys. Rev. B* **13**, 4274 (1976).
- [37] D. Frydel, W. M. Terilla, and K. Burke, *The Journal of Chemical Physics* **112**, 5292 (2000).
- [38] M. Levy and J. P. Perdew, *International journal of quantum chemistry* **49**, 539 (1994).
- [39] M. Levy, *Physical Review A* **43**, 4637 (1991).
- [40] S. Crisostomo, R. Pederson, J. Kozłowski, B. Kalita, A. C. Cancio, K. Datchev, A. Wasserman, S. Song, and K. Burke, *arXiv preprint arXiv:2207.05794* (2022).
- [41] J. P. Perdew and A. Zunger, *Phys. Rev. B* **23**, 5048 (1981).
- [42] C.-O. Almbladh and U. von Barth, *Physical Review B* **31**, 3231 (1985).
- [43] P. Mori-Sánchez, A. J. Cohen, and W. Yang, *Phys. Rev. Lett.* **102**, 066403 (2009).
- [44] A. J. Cohen, P. Mori-Sánchez, and W. Yang, *Chemical reviews* **112**, 289 (2012).
- [45] J. Sun, A. Ruzsinszky, and J. P. Perdew, *Phys. Rev. Lett.* **115**, 036402 (2015).
- [46] A. D. Becke, *The Journal of Chemical Physics* (2022).
- [47] Z. M. Sparrow, B. G. Ernst, T. K. Quady, and R. A. DiStasio Jr, *arXiv preprint arXiv:2109.12560* (2021).
- [48] G. Oliver and J. Perdew, *Phys. Rev. A* **20**, 397 (1979).
- [49] S. Dick and M. Fernandez-Serra, *Physical Review B* **104**, L161109 (2021).
- [50] R. Nagai, R. Akashi, and O. Sugino, *Physical Review Research* **4**, 013106 (2022).
- [51] A. D. Kaplan, S. J. Clark, K. Burke, and J. P. Perdew, *npj Computational Materials* **7**, 1 (2021).
- [52] J. P. Perdew, A. Ruzsinszky, G. I. Csonka, O. A. Vydrov, G. E. Scuseria, L. A. Constantin, X. Zhou, and K. Burke, *Phys. Rev. Lett.* **100**, 136406 (2008).
- [53] M. G. Medvedev, I. S. Bushmarinov, J. Sun, J. P. Perdew, and K. A. Lyssenko, *Science* **355**, 49 (2017).
- [54] B. Kanungo, P. M. Zimmerman, and V. Gavini, *The Journal of Physical Chemistry Letters* **12**, 12012 (2021).
- [55] M. Korth and S. Grimme, *Journal of chemical theory and computation* **5**, 993 (2009).

Supplemental Material

Ryan Pederson*

Department of Physics and Astronomy, University of California, Irvine, CA 92697, USA

Kieron Burke†

Department of Chemistry, University of California, Irvine, CA 92697, USA and
Department of Physics and Astronomy, University of California, Irvine, CA 92697, USA

(Dated: March 6, 2023)

CONTENTS

S1. Gedanken density details	1
S2. computational details	2
S3. Local conditions derivations and relations	2
S4. conventional exchange and correlation partitioning	3
S5. Lieb-Oxford bound for hybrid functionals	3
S6. Local condition for the conjecture: $T_C \leq -E_C$	4
S7. Relation to adiabatic connection	4
S8. Analytical derivation: PBE satisfies the correlation uniform scaling inequality exact condition	5
S9. Readjusting parameters in the SCAN functional	5
S10. exact conditions on atomic system densities	5
S11. assessment of local conditions across available approximations	6
References	12

S1. GEDANKEN DENSITY DETAILS

The Gedanken density used in the main text has the radial form

$$n(r) \propto \begin{cases} C - \frac{A}{\pi} \cos^{-1} \left((1 - \eta) \sin \left(\frac{2\pi r}{T} - \frac{\pi}{2} \right) \right) & r \leq r_p \\ e^{-ar^2 + br + c} & r > r_p \end{cases} \quad (\text{S1})$$

where the density is a dampened triangle wave starting from the origin where $\eta \geq 0$ controls the smoothness of the waveform (where $\eta = 0$ produces a triangle wave), $T > 0$ is the period of the waveform, $N_p \geq 0$ is an integer that controls the number of peaks in the waveform, $A > 0$ is

the amplitude, and $C > 0$ is an offset. At $r_p = (N_p - 3/4)T$, the density transitions to a Gaussian, where a , b , and c are determined by ensuring continuity and first and second derivative continuity. The final gedanken density is then normalized to the desired number of electrons (2 here). Specific variable values used for the gedanken density in the main text are provided in Table S1. In Fig. S1 we plot this gedanken density and its corresponding Kohn-Sham (KS) potential,

$$v_s(r) = \frac{1}{2} \frac{d^2}{dr^2} (r \sqrt{n(r)}). \quad (\text{S2})$$

C	0.2387324146
A	0.1679968844
η	0.05
T	0.3105085788
N_p	5
a	22.0154308155
b	51.4622187780
c	2.2×10^{-14}

TABLE S1. Variable values used in the example gedanken density.

In Table S2, the exchange energy for the gedanken density is given for different exchange approximations, including B88 exchange [1], which was explicitly designed to the correct large r behavior for Coulombic systems. Two-electron densities, such as the gedanken density, follow a tight Lieb-Oxford (LO) bound for exchange, $E_x[n] \geq 1.174 E_x^{\text{LDA}}[n]$ [2]. For the gedanken density, $1.174 E_x^{\text{LDA}} = -0.925$ and we see that PBE and B88 exchange violate this exact condition. On the other hand, SCAN explicitly enforces this tight bound condition and does not violate it.

exact	SCAN	LDA	PBE	B88
-0.832	-0.898	-0.788	-1.062	-1.136

TABLE S2. Exchange energies (in atomic units) computed on the two-electron Gedanken density.

* pedersor@uci.edu

† kieron@uci.edu




FIG. S1. The example Gedanken density considered in the main text and its corresponding KS potential. The potential is continuous everywhere.

S2. COMPUTATIONAL DETAILS

All atomic calculations were performed using the PySCF code [3] with the aug-pcseg-4 basis set. The distribution $g(s)$ was computed using a fermi distribution smoothing with fictitious temperature $T = 0.05$ (following Ref. [4]). For some systems, increased radial grids (up to 500) were used in the Gauss-Chebyshev grid scheme [5] to maintain high fidelity of $g(s)$ at large s values. Further details can be found in our public code [6].

S3. LOCAL CONDITIONS DERIVATIONS AND RELATIONS

We derive several of the local conditions used in the main text which have not been previously reported in the literature to our knowledge. We start with the T_C upper bound exact condition [7]

$$T_C[n_\gamma] \leq -\gamma \left(\frac{\partial E_C[n_\gamma]}{\partial \gamma} \Big|_{\gamma \rightarrow 0} \right) + E_C[n_\gamma], \quad (\text{S3})$$

where [8]

$$T_C[n_\gamma] = \gamma \frac{\partial E_C[n_\gamma]}{\partial \gamma} - E_C[n_\gamma]. \quad (\text{S4})$$

For the semilocal approximations considered here,

$$\tilde{E}_C[n_\gamma] = \gamma \int d^3r n(\mathbf{r}) \epsilon_x^{\text{unif}}[n](\mathbf{r}) \tilde{F}_C(r_s/\gamma, \zeta, s, \alpha, q). \quad (\text{S5})$$

This is due to the fact that the dimensionless quantities ζ , s , α , and q are all scale-invariant, i.e. $\zeta[n_\gamma](\mathbf{r}) =$

$\zeta[n](\gamma\mathbf{r})$, $s[n_\gamma](\mathbf{r}) = s[n](\gamma\mathbf{r})$, and so on. Thus the γ -dependence in the approximation only shows up in the local quantity, $r_s[n_\gamma](\mathbf{r}) = r_s[n](\gamma\mathbf{r})/\gamma$. For compactness, in the following we drop the explicit dependence on scale-invariant quantities. Substituting in Eq. (S5) into Eq. (S3) and enforcing the inequality on the integrands (that is, the local enforcement the exact condition) yields

$$\gamma \frac{\partial \tilde{F}_C(r_s/\gamma)}{\partial \gamma} \geq -\tilde{F}_C(\infty) + \tilde{F}_C(r_s/\gamma). \quad (\text{S6})$$

Using the chain rule and rearranging we obtain the following local condition

$$\frac{\partial \tilde{F}_C}{\partial r_s} \leq \frac{\tilde{F}_C(\infty) - \tilde{F}_C}{r_s}. \quad (\text{S7})$$

The limit $\tilde{F}_C(\infty)$ is the *strictly correlated electron* [9] limit, which is discussed further in Section S7.

The LO bound [10] exact condition reads

$$U_{\text{xc}}[n] \geq C_{\text{LO}} \int d^3r n \epsilon_x^{\text{unif}}[n](\mathbf{r}), \quad (\text{S8})$$

where $U_{\text{xc}}[n] = E_{\text{xc}}[n] - T_C[n]$. For the semilocal approximations considered here, utilizing Eq. (S4) evaluated at $\gamma = 1$ we have

$$\tilde{U}_{\text{xc}}[n] = \int d^3r n(\mathbf{r}) \epsilon_x^{\text{unif}}[n](\mathbf{r}) \left[\tilde{F}_{\text{xc}} + r_s \frac{\partial \tilde{F}_C}{\partial r_s} \right]. \quad (\text{S9})$$

Enforcing the LO bound locally yields the following local condition,

$$\left(1 + r_s \frac{\partial}{\partial r_s} \right) \tilde{F}_{\text{xc}} \leq C_{\text{LO}}. \quad (\text{S10})$$

It is also known that $T_C[n] \geq 0$, and Eq. (S4) yields the following condition

$$\gamma \frac{\partial E_C[n_\gamma]}{\partial \gamma} - E_C[n_\gamma] \geq 0. \quad (\text{S11})$$

However one can show that this condition is automatically satisfied under the *correlation uniform scaling inequality* exact condition [11],

$$\begin{aligned} E_C[n_\gamma] &\geq \gamma E_C[n] & (\gamma > 1), \\ E_C[n_\gamma] &\leq \gamma E_C[n] & (\gamma < 1). \end{aligned} \quad (\text{S12})$$

Let $\gamma > 0$ be arbitrary, $\epsilon > 0$, with $\gamma_+ = 1 + \epsilon/\gamma$ and $\gamma_- = 1 - \epsilon/\gamma$, such that $\epsilon/\gamma \ll 0$ (infinitesimal). From Eq. (S12) we have

$$E_C[n_{\gamma\gamma_+}] - E_C[n_{\gamma\gamma_-}] \geq \gamma_+ E_C[n_\gamma] - \gamma_- E_C[n_\gamma]. \quad (\text{S13})$$

Simplifying yields

$$\gamma \frac{E_C[n_{\gamma+\epsilon}] - E_C[n_{\gamma-\epsilon}]}{2\epsilon} \geq E_C[n_\gamma]. \quad (\text{S14})$$

Applying the definition of a derivative and identifying $T_C[n_\gamma]$ we obtain T_C non-negativity:

$$T_C[n_\gamma] = \gamma \frac{\partial E_C[n_\gamma]}{\partial \gamma} - E_C[n_\gamma] \geq 0. \quad (\text{S15})$$

Enforcing this condition locally yields the following local condition for approximations

$$\frac{\partial \tilde{F}_C(r_s, \zeta, s, \alpha, q)}{\partial r_s} \geq 0, \quad (\text{S16})$$

which is the same one that corresponds to Eq. (S12).

The two inequalities in Eq. (S12) are equivalent. For instance, let $\gamma' < 1$ be arbitrary (but strictly positive) and take $\gamma = 1/\gamma' > 1$. Take $n \mapsto n_{\gamma'}$ in Eq. (S12) and we have $E_C[n_{\gamma'}] \geq \gamma E_C[n]$. Substituting for γ and rearranging we have $E_C[n_{\gamma'}] \leq \gamma' E_C[n]$ with $\gamma' < 1$.

Finally, we note that if the local conditions in Eqs. (S16) and (S7) are satisfied, then we have

$$\tilde{F}_{XC} \leq \tilde{F}_{XC} + r_s \frac{\partial \tilde{F}_C}{\partial r_s} \leq \tilde{F}_X + \tilde{F}_C(\infty). \quad (\text{S17})$$

Since the exchange energy follows the simple scaling relation, $E_X[n_\gamma] = \gamma E_X[n]$, \tilde{F}_X needs to be scale-invariant and thus independent of r_s . The rightmost side of the inequality, $\tilde{F}_X + \tilde{F}_C(\infty)$, is then the maximum value that \tilde{F}_{XC} can take (assuming all other variables besides r_s are fixed). Therefore, if an approximation satisfies Eqs. (S16) and (S7), then the local condition

$$\tilde{F}_{XC} \leq C_{LO}, \quad (\text{S18})$$

which is the standard one that corresponds to the LO bound on E_{XC} , will imply Eq. (S10). In Table 1 in the main text, indeed we see that functionals like PBE, AM05, PBE, and CASE21, which simultaneously satisfy Eqs. (S16), (S7), and (S18), automatically satisfy the LO in Eq. (S8).

We also remark that an enhancement factor \tilde{F}_C that satisfies the conditions in Eqs.(S16), (S7), and (S18) is monotonic and Lipschitz continuous in r_s , i.e., the derivative is bounded, $0 \leq \partial \tilde{F}_C / \partial r_s \leq K$, where K is a finite constant known as the Lipschitz constant. Such a property of the enhancement factor may help assuage possible issues during numerical integration [12]. However, this property is only with respect to the r_s variable and is clearly not sufficient to ensure stability.

S4. CONVENTIONAL EXCHANGE AND CORRELATION PARTITIONING

We define a global hybrid functional approximation as

$$\tilde{E}_{XC}^{\text{hyb}}[n] = \tilde{E}_{XC}[n] + \tilde{a} E_X[n], \quad (\text{S19})$$

where $\tilde{a} > 0$, $E_X[n]$ is the exact exchange energy, and $\tilde{E}_{XC}[n]$ is the remaining semilocal density functional that can be expressed as

$$\tilde{E}_{XC}[n] = \int d^3r n(\mathbf{r}) \tilde{\epsilon}_{XC}(r_s, \zeta, s, \alpha, q). \quad (\text{S20})$$

Note the difference between α and other usual definitions for the mixing parameter in global hybrids.

In hybrid XC functionals and other available approximations, the partitioning of the exchange and correlation may be ambiguous or not defined. In these cases, we use the following conventional partitioning

$$E_C[n] = E_{XC}[n] - \lim_{\gamma \rightarrow \infty} \frac{E_{XC}[n_\gamma]}{\gamma}, \quad (\text{S21})$$

which holds for the exact functional [13]. In global hybrid functionals, the exact exchange contributions cancel out in Eq. (S21), since $E_X[n_\gamma] = \gamma E_X[n]$, and do not contribute to the correlation energy. Here the conventional partitioning for the correlation energy can be expressed as a semilocal density functional with the following correlation energy per electron

$$\tilde{\epsilon}_C[n](\mathbf{r}) = \tilde{\epsilon}_{XC}(r_s, \zeta, s, \alpha, q) - \lim_{\gamma \rightarrow \infty} \frac{\tilde{\epsilon}_{XC}(r_s/\gamma, \zeta, s, \alpha, q)}{\gamma}. \quad (\text{S22})$$

In many cases, the conventional partitioning is consistent with the one defined by the authors of the approximation (if one exists). This agreement occurs whenever an approximation satisfies: $\tilde{E}_X[n_\gamma] = \gamma \tilde{E}_X[n]$ and $\lim_{\gamma \rightarrow \infty} \tilde{E}_C[n_\gamma]/\gamma \rightarrow 0$. The latter is satisfied when $\tilde{E}_C[n_\gamma]$ goes to a finite constant as $\gamma \rightarrow \infty$, but also for approximations of the form

$$\tilde{E}_C[n] = \int d^3r n(\mathbf{r}) \epsilon_C^{\text{unif}}[n] \tilde{G}(\zeta, s, \alpha, q), \quad (\text{S23})$$

where $\epsilon_C^{\text{unif}}[n]$ is a suitable parameterization for the correlation energy per electron of the uniform gas that has logarithmic divergence in the high-density limit (e.g. PW92 [14] or VWN [15]) and $\tilde{G}(\zeta, s, \alpha, q)$ depends only on dimensionless variables. In general, the conventional partitioning we use may differ from the developer's intentions or rationalizations. For instance, approximations that consider a portion of exact exchange to be part of the correlation energy will not be consistent with our conventional partitioning, since such energy contributions will cancel out in Eq. (S21).

Throughout, conventional partitioning is utilized whenever the exchange and correlation partitioning is not available in the Libxc library [16]. In our analysis, we do not consider range-separated hybrid functionals, double-hybrid functionals, or functionals that contain non-local correlation. Therefore, the correlation energy functional is always expressed in the standard semilocal form considered in this work.

S5. LIEB-OXFORD BOUND FOR HYBRID FUNCTIONALS

A global hybrid functional satisfies the XC energy LO bound when

$$\tilde{E}_{XC}[n] \geq C_{LO} E_X^{\text{LDA}}[n] - \tilde{a} E_X[n]. \quad (\text{S24})$$

The exchange energy alone follows a tighter bound

$$E_x[n] \geq C_{\text{LO}}^x E_x^{\text{LDA}}[n], \quad (\text{S25})$$

with coefficient $C_{\text{LO}}^x < C_{\text{LO}}$. Therefore, we can ensure the LO bound with a semilocal functional satisfying

$$\begin{aligned} \tilde{E}_{\text{xc}}[n] &\geq (C_{\text{LO}} - \tilde{a} C_{\text{LO}}^x) E_x^{\text{LDA}}[n] \\ &\geq C_{\text{LO}} E_x^{\text{LDA}}[n] - \tilde{a} E_x[n]. \end{aligned} \quad (\text{S26})$$

Thus, it is sufficient (but not necessary) that $\tilde{E}_{\text{xc}}[n]$ satisfy a LO-like bound with coefficient $C_{\text{LO}} - \tilde{a} C_{\text{LO}}^x$. A larger coefficient could result in a violation of the XC energy LO bound in Eq. (S24). To see this, let $\Delta > 0$ and let n_{LO}^x be a density such that $E_x[n_{\text{LO}}^x] = C_{\text{LO}} E_x^{\text{LDA}}[n_{\text{LO}}^x]$, then $(C_{\text{LO}} - \tilde{a} C_{\text{LO}}^x + \Delta) E_x^{\text{LDA}}[n_{\text{LO}}^x] < C_{\text{LO}} E_x^{\text{LDA}}[n_{\text{LO}}^x] - \tilde{a} E_x[n_{\text{LO}}^x]$. A smaller coefficient would also ensure Eq. (S24), but it would be over-restrictive.

The corresponding local condition is straightforward

$$\tilde{F}_{\text{xc}} \leq C_{\text{LO}} - \tilde{a} C_{\text{LO}}^x. \quad (\text{S27})$$

The corresponding local condition for the LO bound involving $\tilde{U}_{\text{xc}}^{\text{hyb}}[n]$ is found by using the conventional partitioning in Eq. (S22) and applying Eq. (S4) to yield

$$\tilde{F}_{\text{xc}} + r_s \frac{\partial \tilde{F}_{\text{c}}}{\partial r_s} \leq C_{\text{LO}} - \tilde{a} C_{\text{LO}}^x. \quad (\text{S28})$$

In practice, C_{LO} and C_{LO}^x are not known precisely and need to be approximated with proven (but not optimal) bounds. To give the most benefit of the doubt when assessing approximations, we use $C_{\text{LO}}^x = 1.174$ (the conjectured tight exchange LO coefficient [17]) and $C_{\text{LO}} = 2.27$ (which is the same value we use to evaluate non-hybrids).

S6. LOCAL CONDITION FOR THE CONJECTURE: $T_{\text{c}} \leq -E_{\text{c}}$

While unproven, it has been conjectured [8, 11, 18] that

$$T_{\text{c}}[n] \leq -E_{\text{c}}[n] \quad (\text{conjecture}). \quad (\text{S29})$$

One can employ the definition in Eq. (S4) and arrive at the following local condition

$$\frac{\partial \tilde{F}_{\text{c}}}{\partial r_s} \leq \frac{\tilde{F}_{\text{c}}}{r_s} \quad (\text{conjecture}). \quad (\text{S30})$$

In the following Sections S10 and S11 we explore the satisfaction of Eqs. (S29) and (S30), respectively, in approximate functionals.

S7. RELATION TO ADIABATIC CONNECTION

Uniform coordinate scaling is closely related to the adiabatic connection in DFT, which has long been an

illuminating concept for rationalizing and improving density functional approximations [19]. The formalism developed has also revealed many useful exact conditions.

In the adiabatic connection formalism [20–22], we insert a variable coupling constant $\lambda \geq 0$ for Coulomb-interacting electrons with

$$F^\lambda[n] = \min_{\Psi \rightarrow n} \langle \Psi | \hat{T} + \lambda \hat{V}_{\text{ee}} | \Psi \rangle, \quad (\text{S31})$$

where \hat{T} is the usual total kinetic energy operator, \hat{V}_{ee} is the two-body electron-electron repulsion operator, and the minimization is over all antisymmetric wavefunctions that yield the density n . For $\lambda = 1$ and ground-state density $n(\mathbf{r})$, we have our real physical system. Taking $\lambda = 0$, we have the KS system, and $F^{\lambda=0}[n] = T_s[n]$, where $T_s[n]$ is the kinetic energy of the non-interacting KS system with ground-state density $n(\mathbf{r})$. In all cases $\lambda \geq 0$, the ground-state density remains fixed to that of the physical system $n(\mathbf{r})$.

The adiabatic connection is directly coupled to uniform coordinate scaling by

$$F^\lambda[n] = \lambda^2 F[n_{1/\lambda}]. \quad (\text{S32})$$

This relation also extends to any other energy functional component, e.g., $E_{\text{c}}^\lambda[n] = \lambda^2 E_{\text{c}}[n_{1/\lambda}]$. Therefore, exact conditions written in terms of uniform coordinate scaling can be recast in terms of adiabatic connection quantities.

Eq. (S12) is rewritten as

$$\begin{aligned} E_{\text{c}}^\lambda[n] &\geq \lambda E_{\text{c}}[n] \quad (\lambda < 1), \\ E_{\text{c}}^\lambda[n] &\leq \lambda E_{\text{c}}[n] \quad (\lambda > 1). \end{aligned} \quad (\text{S33})$$

Using

$$\frac{\partial E_{\text{c}}[n_{1/\lambda}]}{\partial(1/\lambda)} = -\frac{\partial E_{\text{c}}^\lambda[n]}{\partial \lambda} + 2 \frac{E_{\text{c}}^\lambda[n]}{\lambda} \quad (\text{S34})$$

we recast the exact condition in Eq. (S3) as

$$\frac{\partial E_{\text{c}}^\lambda[n]}{\partial \lambda} \geq \left(-\frac{\partial E_{\text{c}}^\lambda[n]}{\partial \lambda} + 2 \frac{E_{\text{c}}^\lambda[n]}{\lambda} \right) \Big|_{\lambda \rightarrow \infty}, \quad (\text{S35})$$

and applying L'Hospital's rule we obtain

$$\frac{\partial E_{\text{c}}^\lambda[n]}{\partial \lambda} \geq \frac{\partial E_{\text{c}}^\lambda[n]}{\partial \lambda} \Big|_{\lambda \rightarrow \infty}. \quad (\text{S36})$$

This equation can also be rewritten as a simple statement in the *strictly correlated electron* (SCE) [9] limit ($\lambda \rightarrow \infty$),

$$\langle \Psi^\lambda[n] | \hat{V}_{\text{ee}} | \Psi^\lambda[n] \rangle \geq \langle \Psi^{\lambda \rightarrow \infty}[n] | \hat{V}_{\text{ee}} | \Psi^{\lambda \rightarrow \infty}[n] \rangle. \quad (\text{S37})$$

In the SCE limit, the kinetic energy component is subleading and $\Psi^{\lambda \rightarrow \infty}[n]$ minimizes \hat{V}_{ee} , thus Eq. (S37) is clear.

S8. ANALYTICAL DERIVATION: PBE SATISFIES THE CORRELATION UNIFORM SCALING INEQUALITY EXACT CONDITION

The PBE correlation energy functional has the form [23]

$$E_c^{\text{PBE}}[n] = \int d^3r n(\mathbf{r}) \left(\epsilon_c^{\text{PW92}}(r_s, \zeta) + H(r_s, \zeta, t) \right), \quad (\text{S38})$$

where ϵ_c^{PW92} is the PW92 [14] parameterized correlation energy per electron of the uniform gas and $H(r_s, \zeta, t)$ is defined in Ref. [23] along with the dimensionless gradient t . For simplicity, we set all positive constants in H to unity, as the final conclusion will not depend on their specific values. With this,

$$H = \ln \left[1 + t^2 \left[\frac{1 + At^2}{1 + At^2 + A^2 t^4} \right] \right] \geq 0, \quad (\text{S39})$$

where

$$A = [\exp(-\epsilon_c^{\text{PW92}}) - 1]^{-1} \geq 0. \quad (\text{S40})$$

In the following, we show that PBE correlation satisfies Eq. (S16), and thus the correlation uniform scaling inequality exact condition in Eq. (S12). Since PW92 already satisfies Eq. (S16), proving the following condition,

$$\frac{\partial}{\partial r_s} \left(\frac{H(r_s, \zeta, t)}{\epsilon_x^{\text{unif}}(r_s)} \right) \geq 0, \quad (\text{S41})$$

is sufficient to ensure that PBE satisfies the exact condition. To show this, we start with

$$\frac{\partial}{\partial r_s} \left(\frac{H(r_s, \zeta, t)}{\epsilon_x^{\text{unif}}(r_s)} \right) = \frac{(\frac{\partial}{\partial r_s} H)}{\epsilon_x^{\text{unif}}} + \frac{(\frac{\partial}{\partial r_s} \epsilon_x^{\text{unif}}) H}{(\epsilon_x^{\text{unif}})^2}. \quad (\text{S42})$$

Since $\epsilon_x^{\text{unif}} = -(3/4)(3/2\pi)^{2/3} r_s^{-1}$, the second term in Eq. (S42) is positive. Next we evaluate the first term,

$$\frac{\partial H}{\partial r_s} = \frac{1}{1+x} \times \frac{\partial x}{\partial A} \times \frac{\partial A}{\partial \epsilon_c^{\text{PW92}}} \times \frac{\partial \epsilon_c^{\text{PW92}}}{\partial r_s}, \quad (\text{S43})$$

where

$$x = t^2 \left[\frac{1 + At^2}{1 + At^2 + A^2 t^4} \right] \geq 0. \quad (\text{S44})$$

The intermediate derivatives are derived analytically:

$$\frac{\partial x}{\partial A} = \frac{-t^4 A (t^2 A + 2)}{(t^4 A^2 + t^2 A + 1)^2} \leq 0, \quad (\text{S45})$$

$$\frac{\partial A}{\partial \epsilon_c^{\text{PW92}}} = \frac{\exp(-\epsilon_c^{\text{PW92}})}{(\exp(-\epsilon_c^{\text{PW92}}) - 1)^2} \geq 0, \quad (\text{S46})$$

and

$$\frac{\partial \epsilon_c^{\text{PW92}}}{\partial r_s} \geq 0. \quad (\text{S47})$$

Therefore,

$$\frac{\partial H}{\partial r_s} \leq 0, \quad (\text{S48})$$

and the first term in Eq. (S42) is also positive and thus the exact condition is satisfied.

S9. READJUSTING PARAMETERS IN THE SCAN FUNCTIONAL

While the local condition in Eq. (S16) is satisfied in the published SCAN functional, reasonable adjustments of the parameters designated for the fitting of the ‘‘appropriate norms’’ can result in violations of the local condition. Specifically, we adjust the $b_{1c} = 0.02858$ parameter in SCAN, which was fit to match the correlation energy of the $Z \rightarrow \infty$ limit of two-electron ions (one of the five appropriate norms in SCAN, see the supplementary material in Ref. [24]). By evaluating SCAN analytically, we find that using $b_{1c} > 0.2$ results in violations of Eq. (S16). Under this single parameter modification, the other exact conditions that SCAN satisfies are still satisfied by construction. Therefore, by virtue of fitting to various appropriate norms, the SCAN functional satisfies more exact conditions than were explicitly enforced.

S10. EXACT CONDITIONS ON ATOMIC SYSTEM DENSITIES

In this assessment, we first calculate Hartree-Fock (HF) densities and orbitals for neutral atoms H-Ar and their cations. The fixed HF densities and orbitals are then used to evaluate the energies (non-self-consistently) from different DFT approximations. HF densities are used because they provide high quality densities and an equal footing across different approximations. We also performed separate calculations using self-consistent densities and observed marginal differences. The absolute errors from the experimental ionization energies are provided in Fig. S2. In addition to established approximations, we also test a very simple modified B3LYP (MOD-B3LYP) which satisfies the correlation exact conditions discussed for any density,

$$\epsilon_c^{\text{MOD-B3LYP}}(r_s, \zeta, s) = \Theta(s - 1.82) \epsilon_c^{\text{B3LYP}}(r_s, \zeta, s), \quad (\text{S49})$$

where $\epsilon_c^{\text{B3LYP}} = 0.405\epsilon_c^{\text{LYP}} + 0.095\epsilon_c^{\text{VWN5}}$ [25]. The step function eliminates the local condition violations found in Table 1 of the main text. As argued in the main text, energy contributions from such large s values are less relevant in Coulombic systems, and indeed in Fig. S2 we see that B3LYP and MOD-B3LYP have MAEs that differ only by 0.1 kcal/mol. Our modified functional is constructed for demonstration purposes only.

For each HF density (a total of 35 atomic systems), we scale the density n_γ with $\gamma \in [0.01, 2]$ (50 evenly spaced values) and evaluate whether an exact condition is satisfied. Indeed, in Fig. S3 we see that the exact conditions tested are all satisfied within our set of functional approximations and atomic systems. We also test the conjecture $T_c \leq -E_c$ and find instances of violation for PBE and M08-HX.




FIG. S2. Ionization energy errors for atomic systems H-Ar. All approximations are evaluated on HF densities and orbitals.




FIG. S3. The number of HF atomic densities tested which satisfy a given exact condition. A total of 35 atomic systems are tested: H-Ar and their cations.

S11. ASSESSMENT OF LOCAL CONDITIONS ACROSS AVAILABLE APPROXIMATIONS

We utilize an exhaustive grid search to determine whether local conditions are satisfied for a given approximation. For GGAs, we consider 10000 evenly spaced values of $r_s \in [0.0001, 5]$, 500 evenly spaced values of $s \in [0, 5]$, and 100 evenly spaced values of $\zeta \in [0, 1]$. For MGGAs, we consider 5000 evenly spaced values of $r_s \in [0.0001, 5]$, 100 evenly spaced values of $s \in [0, 5]$, 20 evenly spaced values of $\zeta \in [0, 1]$, and 100 evenly spaced values of $\alpha \in [0, 5]$ or $q \in [-10, 10]$. The number of values checked per variable is less in MGGAs to alleviate computational effort due to the combinatorial nature of the exhaustive search.

In determining whether local conditions are satisfied, a reasonable tolerance threshold of at most ± 0.001 is employed to approximately account for numerical errors arising from the numerical precision used and the finite difference method used to calculate numerical

derivatives (further details can be found in our public code [6]). However, the numerical errors introduced are not guaranteed to be within the tolerances used.

In the following tables below, we report the fraction of local condition violations found in our exhaustive search. That is, we divide the number of violations found by the total number of permutations considered in the extensive grid search parameter space. If we find 0 such violations, then we conclude that the corresponding exact condition is satisfied for any reasonable density.

TABLE S3. GGA functionals: numerical assessment of corresponding local conditions.

	$E_C[n]$ non-positivity	$E_C[n_\gamma]$ uniform scaling inequality	$T_C[n]$ upper bound	$U_C(\lambda)$ monotonicity	LO extension to E_{XC}	LO	conjecture: $T_c \leq -E_c$
ACGGAP [26, 27]	0	0	0	0	—	—	0.414
ACGGA [26, 27]	0	0	0	0	—	—	0
AM05 [28, 29]	0	0	0	0	0	0	0
APBE [30]	0	0	0	0	0	0	0.004
B97-D [31]	0.632	0.503	0.559	0.608	0.268	0.183	0.633
B97-GGA1 [32]	0.636	0.514	0.557	0.612	0.390	0.317	0.639
BEEFVDW [33]	0	0	0	0	0.003	0.013	0
BMK [34]	0.627	0.304	0.648	0.621	—	—	0.616
CCDF [35]	0	0	0	0	—	—	0
CHACHIYO [36]	0	0	0.044	0	0.217	0.217	0.010
CS1 [37, 38]	0.604	0.204	0.528	0.530	—	—	0.601
EDF1 [39]	0.605	0.245	0.002	0.231	0.162	0.203	0.527
FT97 [40, 41]	0	0	1e-05	0.003	—	—	0
GAM [42]	0.598	0.459	0.560	0.578	0.145	0.083	0.596
GAPC [43]	0.004	0.011	2e-04	0.005	—	—	0.015
GAPLOC [43]	4e-04	2e-04	2e-04	0.005	—	—	0.033
HCTH-120 [44]	0.495	0.310	0.327	0.450	0.065	0.061	0.507
HCTH-147 [44]	0.467	0.290	0.298	0.422	0.113	0.093	0.478
HCTH-407P [45]	0.536	0.445	0.428	0.508	0.105	0.075	0.543
HCTH-407 [46]	0.481	0.382	0.365	0.450	0.112	0.079	0.489
HCTH-93 [47]	0.435	0.196	0.258	0.386	0.266	0.237	0.446
HCTH-A [47]	0.493	0.289	0.355	0.454	0	0	0.501
HCTH-P14 [48]	0	0	0	0	0	0.029	0
HCTH-P76 [48]	0.986	0.929	0.999	0.991	0.011	0.005	0.978
HLE16 [49]	0.481	0.305	0.364	0.447	0.477	0.473	0.487
HYB-TAU-HCTH [50]	0.615	0.439	0.520	0.585	—	—	0.620
KT1 [51]	0.791	0.434	0.096	0.169	0.153	0.076	0.664
KT2 [51]	0.832	0.477	0.106	0.177	0.156	0.077	0.686
KT3 [52]	0.862	0.461	0.120	0.192	0.164	0.077	0.678
LM [53, 54]	0	0.119	0.464	0.384	—	—	0
LYPR [55]	0.320	0.113	0.801	0.590	—	—	0.438
LYP [56, 57]	0.576	0.218	0.003	0.203	—	—	0.511
MGGAC [58]	0	0	0	0	—	—	0.007
MOHLYP2 [59]	0.576	0.174	0.002	0.193	0.340	0.337	0.509
MOHLYP [60]	0.243	0.092	0	0.092	0.048	0.096	0.328
MPWLYP1W [61]	0.500	0.190	4e-07	0.168	0.003	0.004	0.474
N12 [62]	0	0	0	0	0.150	0.170	0
NCAP [63]	0.455	0.299	0.029	0.277	0.231	0.207	0.403
OBLYP-D [64]	0.595	0.246	0.002	0.243	0.019	0.020	0.505
OP-B88 [65]	0	2e-04	6e-04	0.002	—	—	7e-04
OP-G96 [65, 66]	0	2e-04	6e-04	0.002	—	—	7e-04
OP-PBE [65, 66]	0	2e-04	6e-04	0.002	—	—	7e-04
OP-PW91 [65, 66]	0	0.001	0.001	0.002	—	—	0.001
OP-XALPHA [65, 66]	0	2e-04	6e-04	0.002	—	—	7e-04
OPBE-D [64]	0	0	0	0	0.009	0.009	0.006
OPTC [67]	3e-06	0	0	0	—	—	0
OPWLYP-D [64]	0.596	0.247	7e-04	0.240	0.019	0.022	0.507
P86-FT [68]	0.454	0.298	0.027	0.276	—	—	0.402
P86VWN-FT [68]	0.447	0.297	0.026	0.275	—	—	0.389
P86VWN [68]	0.447	0.297	0.026	0.275	—	—	0.389
P86 [68]	0.454	0.298	0.027	0.276	—	—	0.403
PBE-JRGX [69]	0	0	0	0	—	—	0.006
PBE-MOL [70]	0	0	0	0	0	0	0.004
PBE-SOL [71]	0	0	0	0	0	0	0.006
PBE-VWN [72–74]	0	0	0	0	—	—	6e-04
PBE1W [61]	0	0	0	0	0	0	0
PBEFE [75]	0	0	0	0	0	0	0.006
PBEINT [76]	0	0	0	0	0	0	0.005
PBELOC [77]	0	0	0	0.003	—	—	0.271
PBELYP1W [61]	0.414	0.157	0	0.141	0	0	0.427
PBE [73, 74]	0	0	0	0	0	0	0.005

TABLE S4. GGA functionals: numerical assessment of corresponding local conditions.

	$E_C[n]$ non-positivity	$E_C[n_\gamma]$ uniform scaling inequality	$T_C[n]$ upper bound	$U_C(\lambda)$ monotonicity	LO extension to E_{XC}	LO	conjecture: $T_c \leq -E_c$
PW91 [78–80]	3e-06	0	0	0	0	0	0
Q2D [81]	0	0.041	0.012	0.032	0	0	0.002
REGTPSS [82]	0	0	0	0	—	—	0.406
REVTCA [83]	0	0	0.003	0.051	—	—	0.024
RGE2 [84]	0	0	0	0	0	0	0.005
SCAN-E0 [85]	0	0	0	0	—	—	0
SG4 [86]	0	0.048	0.327	0.385	8e-04	0.007	0.050
SOGGA11 [87]	0	0.003	0.064	0.229	0	1e-04	0.002
SPBE [88]	0	0	0	0	—	—	0
TAU-HCTH [50]	0.595	0.491	0.492	0.567	—	—	0.603
TCA [89]	0	0	0	0	—	—	0
TH-FCFO [90]	0.223	0.772	0.233	0.237	0.787	0.228	0.226
TH-FCO [90]	0.200	0.795	0.211	0.214	0.781	0.205	0.203
TH-FC [90]	0.988	0.009	0.994	0.996	0.822	0.990	0.989
TH-FL [90]	0	1.000	0	0	0.498	0	0
TH1 [91]	0.215	0.780	0.224	0.225	0.295	0.220	0.218
TH2 [92]	0.061	0.935	0.070	0.070	0.323	0.065	0.063
TH3 [93]	0.217	0.781	0.217	0.219	0.284	0.218	0.218
TH4 [93]	0.103	0.894	0.106	0.106	0.238	0.105	0.105
TM-LYP [94]	0.575	0.209	0.122	0.173	—	—	0.565
TM-PBE [94]	0	0	0	0	—	—	0.509
W94 [95]	0	0	0	3e-05	—	—	0
W10 [96]	0.614	0.004	0.002	0.014	—	—	0.603
WI [96]	0.900	0.008	0.008	0.023	—	—	0.896
WL [97]	0.590	0.166	0.595	0.581	—	—	0.377
XLYP [98]	0.576	0.218	0.003	0.203	0.013	0.019	0.511
XPBE [99]	0	0	0	0	0	0	0
ZPBEINT [100]	0	0.020	0.343	0.299	—	—	2e-04
ZPBESOL [100]	0	0.013	0.375	0.263	—	—	2e-04
ZVPBEINT [101]	0	0.093	0.273	0.226	—	—	0.001
ZVPBELOC [102]	0	3e-04	0.199	0.112	—	—	0.075
ZVPBESOL [101]	0	0.080	0.299	0.212	—	—	0.001

TABLE S5. MGGA functionals: numerical assessment of corresponding local conditions.

	$E_c[n]$ non-positivity	$E_c[n_\gamma]$ uniform scaling inequality	$T_c[n]$ upper bound	$U_c(\lambda)$ monotonicity	LO extension to E_{xc}	LO	conjecture: $T_c \leq -E_c$
B88 [103]	0	0	0	0	—	—	0
B94 [104]	0	0	0	0	—	—	0
BC95 [105]	0	0	0	0	—	—	0
CC06 [106]	0	0	0	0	0	0	0
CS [56, 107]	0.352	0.166	0.252	0.275	—	—	0.481
HLE17 [108]	0	0	0	0	0.092	0.093	0
HLTAPW [109]	0	0	0	0	—	—	0
KCISK [110–114]	0	0	0	0.012	—	—	0.041
KCIS [110–114]	0	0	0	0	—	—	0
LP90 [115]	0	0.963	0.998	0.990	0	0	0
M06-L [116, 117]	0.700	0.661	0.705	0.696	0.228	0.181	0.698
M11-L [118]	0.385	0.194	0.004	0.153	0.425	0.456	0.478
MN12-L [119]	0.424	0.266	0.022	0.227	0.048	0.086	0.506
MN15-L [120]	0.462	0.184	3e-04	0.156	2e-04	0.006	0.594
OTPSS-D [64]	0	0	0	0	0	0	0.007
PKZB [121]	0	0	0	0	0	0	0.006
R2SCANL [122–124]	0	0	0	0	0	0	0
R2SCAN [123, 124]	0	0	0	0	0	0	0
REVM06-L [125]	0.777	0.702	0.835	0.783	5e-05	7e-05	0.767
REVSCAN [126]	0.212	0.063	0	0.084	0	0	0.288
REVTM [127]	0	0	0	0	0	0	0.329
REVPSS [82, 128]	0	0	0	0	0	0	0.424
RSCAN [129]	0	0.008	0.101	0.160	0	0	0
SCANL [85, 130, 131]	0	0	0	0	0	0	0
SCAN [85]	0	0	0	0	0	0	0
TM [132]	0	0	0	0	0	0	7e-04
TPSSLOC [77]	0	0	0	0.003	—	—	0.280
TPSSLYP1W [61]	0.408	0.179	0	0.138	0	0	0.420
TPSS [133, 134]	0	0	0	0	0	0	0.008
VSXC [135]	0.298	0.084	0.190	0.260	—	—	0.294
ZLP [136]	0	0.943	0.998	0.981	0.343	0.285	0

TABLE S6. Hybrid GGA functionals: numerical assessment of corresponding local conditions.

	$E_C[n]$ non-positivity	$E_C[n_\gamma]$ uniform scaling inequality	$T_C[n]$ upper bound	$U_C(\lambda)$ monotonicity	LO extension to E_{XC}	LO	conjecture: $T_c \leq -E_c$
APBE0 [102]	0	0	0	0	0	0	0.004
APF [137]	0	0	0	0	0	0	0
B1LYP [138]	0.576	0.218	0.003	0.203	0	0	0.511
B1PW91 [138]	0	0	0	0	0	0	0
B1WC [139]	0	0	0	0	0	0	0.005
B3LYP-MCM1 [140]	0.753	0.282	0.681	0.455	0	0	0.590
B3LYP-MCM2 [140]	0.558	0.210	4e-04	0.187	0	0	0.504
B3LYP3 [141]	0.457	0.174	0	0.154	0	0	0.451
B3LYP5 [141]	0.457	0.174	0	0.154	0	0	0.451
B3LYPS [142]	0.414	0.159	0	0.137	0	5e-05	0.442
B3LYP [141]	0.414	0.159	0	0.137	0	8e-05	0.442
B3P86 [143]	0.074	0.141	0	0.209	0	0	0.265
B3PW91 [144]	0	0	0	0	0	0	0
B5050LYP [145]	0.457	0.174	0	0.154	0	0	0.451
B97-1P [32]	0.666	0.463	0.634	0.649	0.056	0.042	0.663
B97-1 [47]	0.698	0.476	0.686	0.687	0.002	0.001	0.693
B97-2 [146]	0.665	0.479	0.605	0.643	0.065	0.048	0.666
B97-3 [147]	0.612	0.464	0.601	0.601	0.089	0.050	0.608
B97-K [34]	0.330	0.004	0.520	0.342	0	0	0.291
B97 [148]	0.645	0.359	0.612	0.626	0.003	0.003	0.641
BHANDHLYP [143, 149]	0.576	0.218	0.003	0.203	0	0	0.511
BHANDH [143, 149]	0.576	0.218	0.003	0.203	0	0	0.511
BLYP35 [150, 151]	0.576	0.218	0.003	0.203	0	0	0.511
CAPO [152]	0	0	0	0	0.080	0.081	0.005
CASE21 [153]	0	0	0	0	0	0	0.259
EDF2 [154]	0.362	0.132	0	0.125	0	0	0.398
HAPBE [102]	0	0	0	0	0	0	0.012
HFLYP [56, 57]	0.576	0.218	0.003	0.203	0	0	0.511
HPBEINT [155]	0	0	0	0	0	0	0.005
KMLYP [156]	0.121	0.052	0	0.054	0	0	0.279
MB3LYP-RC04 [157]	0.359	0.143	0.001	0.126	0	1e-04	0.375
MPW1K [158]	0	0	0	0	0	0	0
MPW1LYP [159]	0.576	0.218	0.003	0.203	0	0	0.511
MPW1PBE [159]	0	0	0	0	0	0	0.005
MPW1PW [159]	0	0	0	0	0	0	0
MPW3LYP [160]	0.462	0.178	0	0.153	0	0	0.465
MPW3PW [159]	0	0	0	0	0	0	0
MPWLYP1M [60]	0.576	0.218	0.003	0.203	3e-04	3e-04	0.511
O3LYP [67, 161]	0.457	0.174	0	0.154	0.017	0.028	0.451
PBE-2X [162]	0	0	0	0	0	0	0.005
PBE-MOLO [70]	0	0	0	0	0	0	0.004
PBE-MOLB0 [70]	0	0	0	0	0	0	0.005
PBE-SOLO [70]	0	0	0	0	0	0	0.006
PBE0-13 [163]	0	0	0	0	0	0	0.005
PBE38 [164]	0	0	0	0	0	0	0.005
PBE50 [165]	0	0	0	0	0	0	0.005
PBEB0 [70]	0	0	0	0	0	0	0.005
PBEH [166, 167]	0	0	0	0	0	0	0.005
QTP17 [168]	0.407	0.156	0	0.135	0	0	0.439
REVB3LYP [169]	0.438	0.168	0	0.145	0	0	0.453
SB98-1A [170]	0.803	0.435	0.903	0.809	0	0	0.786
SB98-1B [170]	0.267	0.049	0.055	0.215	0.013	0.019	0.278
SB98-1C [170]	0.635	0.363	0.593	0.615	0.004	0.004	0.633
SB98-2A [170]	0	0	0	0	0.002	0.012	0
SB98-2B [170]	0.478	0.180	0.327	0.432	2e-05	4e-04	0.484
SB98-2C [170]	0.632	0.371	0.586	0.611	0.001	0.002	0.629
SOGGA11-X [171]	0	0.042	0.176	0.169	0	0.003	0
WC04 [172]	0	0	0	0	0.123	0.240	0
WP04 [172]	0	0	0	0	0.349	0.468	0
X3LYP [98]	0.462	0.178	0	0.153	0	0	0.465

TABLE S7. Hybrid MGGA functionals: numerical assessment of corresponding local conditions.

	$E_C[n]$ non-positivity	$E_C[n_\gamma]$ uniform scaling inequality	$T_C[n]$ upper bound	$U_C(\lambda)$ monotonicity	LO extension to E_{XC}	LO	conjecture: $T_c \leq -E_c$
B0KCIS [114]	0	0	0	0	0	0	0
B86B95 [105]	0	0	0	0	0	9e-04	0
B88B95 [105]	0	0	0	0	0	9e-04	0
B98 [173]	0.202	0.004	0.145	0.119	0	0	0.172
BB1K [174]	0	0	0	0	0	1e-03	0
BR3P86 [175]	0.290	0.336	0.005	0.247	0	6e-04	0.305
DLDF [176]	0.543	0.455	0.523	0.532	0.298	0.241	0.540
EDMGGAH [177]	0.351	0.166	0.254	0.276	0.015	0.013	0.481
M05-2X [178]	0.745	0.696	0.760	0.743	0.070	0.042	0.741
M05 [179]	0.748	0.559	0.767	0.729	0.075	0.073	0.731
M06-2X [117]	0.681	0.613	0.720	0.682	0.058	0.036	0.674
M06-HF [180]	0.298	0.232	0.382	0.313	0.049	0.105	0.284
M06 [117]	0.419	0.229	0.257	0.375	0.329	0.330	0.423
M08-HX [181]	0.039	0.008	0	0.007	0.070	0.259	0.136
M08-SO [181]	0.079	0.019	0	0.015	0.149	0.260	0.196
MN15 [182]	0.785	0.612	0.396	0.631	0.288	0.276	0.860
MPW1B95 [160]	0	0	0	0	0	9e-04	0
MPW1KCIS [183]	0	0	0	0	0	0	0
MPWB1K [160]	0	0	0	0	0	1e-03	0
MPWKCIS1K [183]	0	0	0	0	0	0	0
PBE1KCIS [184]	0	0	0	0	0	0	0
PW6B95 [185]	0	0	0	0	0	9e-04	0
PW86B95 [105]	0	0	0	0	0	9e-04	0
PWB6K [185]	0	0	0	0	0	1e-03	0
REVM06 [186]	0.676	0.591	0.685	0.670	0.003	0.002	0.670
REVTPSSH [187]	0	0	0	0	0	0	0.424
TPSS0 [188]	0	0	0	0	0	0	0.008
TPSS1KCIS [189]	0	0	0	0	0	0	0
TPSSH [190]	0	0	0	0	0	0	0.008
X1B95 [160]	0	0	0	0	0	0	0
XB1K [160]	0	0	0	0	0	4e-05	0

- [1] A. D. Becke, Physical review A **38**, 3098 (1988).
- [2] J. P. Perdew, A. Ruzsinszky, J. Sun, and K. Burke, The Journal of Chemical Physics **140**, 18A533 (2014).
- [3] Q. Sun, X. Zhang, S. Banerjee, P. Bao, M. Barbry, N. S. Blunt, N. A. Bogdanov, G. H. Booth, J. Chen, Z.-H. Cui, *et al.*, The Journal of chemical physics **153**, 024109 (2020).
- [4] A. Zupan, J. P. Perdew, K. Burke, and M. Causà, International journal of quantum chemistry **61**, 835 (1997).
- [5] M. Krack and A. M. Köster, The Journal of chemical physics **108**, 3226 (1998).
- [6] https://github.com/pedersor/dft_exconditions (2023).
- [7] M. Levy and J. P. Perdew, International journal of quantum chemistry **49**, 539 (1994).
- [8] D. Frydel, W. M. Terilla, and K. Burke, The Journal of Chemical Physics **112**, 5292 (2000).
- [9] M. Seidl, P. Gori-Giorgi, and A. Savin, Physical Review A **75**, 042511 (2007).
- [10] E. H. Lieb and S. Oxford, International Journal of Quantum Chemistry **19**, 427 (1981).
- [11] M. Levy and J. P. Perdew, Physical Review A **32**, 2010 (1985).
- [12] S. Lehtola and M. A. Marques, arXiv preprint arXiv:2206.14062 (2022).
- [13] M. Levy and J. P. Perdew, Physical Review B **48**, 11638 (1993).
- [14] J. P. Perdew and Y. Wang, Physical review B **45**, 13244 (1992).
- [15] S. H. Vosko, L. Wilk, and M. Nusair, Canadian Journal of physics **58**, 1200 (1980).
- [16] S. Lehtola, C. Steigemann, M. J. Oliveira, and M. A. Marques, SoftwareX **7**, 1 (2018).
- [17] J. P. Perdew and J. Sun, arXiv preprint arXiv:2206.09974 (2022).
- [18] S. Crisostomo, R. Pederson, J. Kozłowski, B. Kalita, A. C. Cancio, K. Datchev, A. Wasserman, S. Song, and K. Burke, arXiv preprint arXiv:2207.05794 (2022).
- [19] J. P. Perdew, M. Ernzerhof, and K. Burke, The Journal of Chemical Physics **105**, 9982 (1996).
- [20] J. Harris and R. Jones, J. Phys. F **4**, 1170 (1974).
- [21] D. Langreth and J. Perdew, Solid State Commun. **17**, 1425 (1975).
- [22] O. Gunnarsson and B. Lundqvist, Phys. Rev. B **13**, 4274 (1976).
- [23] J. P. Perdew, K. Burke, and M. Ernzerhof, Phys. Rev. Lett. **77**, 3865 (1996), *ibid.* **78**, 1396(E) (1997).
- [24] J. Sun, A. Ruzsinszky, and J. P. Perdew, Phys. Rev. Lett. **115**, 036402 (2015).
- [25] P. J. Stephens, F. J. Devlin, C. F. Chabalowski, and M. J. Frisch, The Journal of physical chemistry **98**, 11623 (1994).
- [26] A. Cancio, G. P. Chen, B. T. Krull, and K. Burke, J. Chem. Phys. **149**, 084116 (2018).
- [27] K. Burke, A. Cancio, T. Gould, and S. Pittalis, ArXiv e-prints (2014), arXiv:1409.4834 [cond-mat.mtrl-sci].
- [28] R. Armiento and A. E. Mattsson, Phys. Rev. B **72**, 085108 (2005).
- [29] A. E. Mattsson, R. Armiento, J. Paier, G. Kresse, J. M. Wills, and T. R. Mattsson, J. Chem. Phys. **128**, 084714 (2008).
- [30] L. A. Constantin, E. Fabiano, S. Laricchia, and F. Della Sala, Phys. Rev. Lett. **106**, 186406 (2011).
- [31] S. Grimme, J. Comput. Chem. **27**, 1787 (2006).
- [32] A. J. Cohen and N. C. Handy, Chem. Phys. Lett. **316**, 160 (2000).
- [33] J. Wellendorff, K. T. Lundgaard, A. Møgelhøj, V. Petzold, D. D. Landis, J. K. Nørskov, T. Bligaard, and K. W. Jacobsen, Phys. Rev. B **85**, 235149 (2012).
- [34] A. D. Boese and J. M. L. Martin, J. Chem. Phys. **121**, 3405 (2004).
- [35] J. T. Margraf, C. Kunkel, and K. Reuter, J. Chem. Phys. **150**, 244116 (2019).
- [36] T. Chachiyo and H. Chachiyo, Comput. Theor. Chem. **1172**, 112669 (2020).
- [37] N. C. Handy and A. J. Cohen, J. Chem. Phys. **116**, 5411 (2002).
- [38] E. I. Proynov and A. J. Thakkar, Int. J. Quantum Chem. **106**, 436 (2006).
- [39] R. D. Adamson, P. M. W. Gill, and J. A. Pople, Chem. Phys. Lett. **284**, 6 (1998).
- [40] M. Filatov and W. Thiel, Int. J. Quantum Chem. **62**, 603 (1997).
- [41] M. Filatov and W. Thiel, Mol. Phys. **91**, 847 (1997).
- [42] H. S. Yu, W. Zhang, P. Verma, X. He, and D. G. Truhlar, Phys. Chem. Chem. Phys. **17**, 12146 (2015).
- [43] E. Fabiano, P. E. Trevisanutto, A. Terentjev, and L. A. Constantin, J. Chem. Theory Comput. **10**, 2016 (2014).
- [44] A. D. Boese, N. L. Doltsinis, N. C. Handy, and M. Sprik, J. Chem. Phys. **112**, 1670 (2000).
- [45] A. D. Boese, A. Chandra, J. M. L. Martin, and D. Marx, J. Chem. Phys. **119**, 5965 (2003).
- [46] A. D. Boese and N. C. Handy, J. Chem. Phys. **114**, 5497 (2001).
- [47] F. A. Hamprecht, A. J. Cohen, D. J. Tozer, and N. C. Handy, J. Chem. Phys. **109**, 6264 (1998).
- [48] G. Menconi, P. J. Wilson, and D. J. Tozer, J. Chem. Phys. **114**, 3958 (2001).
- [49] P. Verma and D. G. Truhlar, J. Phys. Chem. Lett. **8**, 380 (2017).
- [50] A. D. Boese and N. C. Handy, J. Chem. Phys. **116**, 9559 (2002).
- [51] T. W. Keal and D. J. Tozer, J. Chem. Phys. **119**, 3015 (2003).
- [52] T. W. Keal and D. J. Tozer, J. Chem. Phys. **121**, 5654 (2004).
- [53] D. C. Langreth and M. J. Mehl, Phys. Rev. Lett. **47**, 446 (1981).
- [54] C. D. Hu and D. C. Langreth, Phys. Scr. **32**, 391 (1985).
- [55] W. Ai, W.-H. Fang, and N. Q. Su, J. Phys. Chem. Lett. **12**, 1207 (2021).
- [56] C. Lee, W. Yang, and R. G. Parr, Phys. Rev. B **37**, 785 (1988).
- [57] B. Miehlich, A. Savin, H. Stoll, and H. Preuss, Chem. Phys. Lett. **157**, 200 (1989).
- [58] B. Patra, S. Jana, L. A. Constantin, and P. Samal, Phys. Rev. B **100**, 155140 (2019).
- [59] J. Zheng, Y. Zhao, and D. G. Truhlar, J. Chem. Theory Comput. **5**, 808 (2009).
- [60] N. E. Schultz, Y. Zhao, and D. G. Truhlar, J. Phys. Chem. A **109**, 11127 (2005).

- [61] E. E. Dahlke and D. G. Truhlar, *J. Phys. Chem. B* **109**, 15677 (2005).
- [62] R. Peverati and D. G. Truhlar, *J. Chem. Theory Comput.* **8**, 2310 (2012).
- [63] J. Carmona-Espíndola, J. L. Gázquez, A. Vela, and S. B. Trickey, *J. Chem. Theory Comput.* **15**, 303 (2019).
- [64] L. Goerigk and S. Grimme, *J. Chem. Theory Comput.* **6**, 107 (2010).
- [65] T. Tsuneda, T. Suzumura, and K. Hirao, *J. Chem. Phys.* **110**, 10664 (1999).
- [66] T. Tsuneda, T. Suzumura, and K. Hirao, *J. Chem. Phys.* **111**, 5656 (1999).
- [67] A. J. Cohen and N. C. Handy, *Mol. Phys.* **99**, 607 (2001).
- [68] J. P. Perdew, *Phys. Rev. B* **33**, 8822 (1986).
- [69] L. S. Pedroza, A. J. R. da Silva, and K. Capelle, *Phys. Rev. B* **79**, 201106 (2009).
- [70] J. M. del Campo, J. L. Gázquez, S. B. Trickey, and A. Vela, *J. Chem. Phys.* **136**, 104108 (2012).
- [71] J. P. Perdew, A. Ruzsinszky, G. I. Csonka, O. A. Vydrov, G. E. Scuseria, L. A. Constantin, X. Zhou, and K. Burke, *Phys. Rev. Lett.* **100**, 136406 (2008).
- [72] E. Kraisler, G. Makov, and I. Kelson, *Phys. Rev. A* **82**, 042516 (2010).
- [73] J. P. Perdew, K. Burke, and M. Ernzerhof, *Phys. Rev. Lett.* **77**, 3865 (1996).
- [74] J. P. Perdew, K. Burke, and M. Ernzerhof, *Phys. Rev. Lett.* **78**, 1396 (1997).
- [75] R. Sarmiento-Pérez, S. Botti, and M. A. L. Marques, *J. Chem. Theory Comput.* **11**, 3844 (2015).
- [76] E. Fabiano, L. A. Constantin, and F. Della Sala, *Phys. Rev. B* **82**, 113104 (2010).
- [77] L. A. Constantin, E. Fabiano, and F. Della Sala, *Phys. Rev. B* **86**, 035130 (2012).
- [78] J. P. Perdew, in *Proceedings of the 75. WE-Heraeus-Seminar and 21st Annual International Symposium on Electronic Structure of Solids*, edited by P. Ziesche and H. Eschrig (Akademie Verlag, Berlin, 1991) p. 11.
- [79] J. P. Perdew, J. A. Chevary, S. H. Vosko, K. A. Jackson, M. R. Pederson, D. J. Singh, and C. Fiolhais, *Phys. Rev. B* **46**, 6671 (1992).
- [80] J. P. Perdew, J. A. Chevary, S. H. Vosko, K. A. Jackson, M. R. Pederson, D. J. Singh, and C. Fiolhais, *Phys. Rev. B* **48**, 4978 (1993).
- [81] L. Chiodo, L. A. Constantin, E. Fabiano, and F. Della Sala, *Phys. Rev. Lett.* **108**, 126402 (2012).
- [82] J. P. Perdew, A. Ruzsinszky, G. I. Csonka, L. A. Constantin, and J. Sun, *Phys. Rev. Lett.* **103**, 026403 (2009).
- [83] V. Tognetti, P. Cortona, and C. Adamo, *Chem. Phys. Lett.* **460**, 536 (2008).
- [84] A. Ruzsinszky, G. I. Csonka, and G. E. Scuseria, *J. Chem. Theory Comput.* **5**, 763 (2009).
- [85] J. Sun, A. Ruzsinszky, and J. P. Perdew, *Phys. Rev. Lett.* **115**, 036402 (2015).
- [86] L. A. Constantin, A. Terentjevs, F. Della Sala, P. Cortona, and E. Fabiano, *Phys. Rev. B* **93**, 045126 (2016).
- [87] R. Peverati, Y. Zhao, and D. G. Truhlar, *J. Phys. Chem. Lett.* **2**, 1991 (2011).
- [88] M. Swart, M. Solá, and F. M. Bickelhaupt, *J. Chem. Phys.* **131**, 094103 (2009).
- [89] V. Tognetti, P. Cortona, and C. Adamo, *J. Chem. Phys.* **128**, 034101 (2008).
- [90] D. J. Tozer, N. C. Handy, and W. H. Green, *Chem. Phys. Lett.* **273**, 183 (1997).
- [91] D. J. Tozer and N. C. Handy, *J. Chem. Phys.* **108**, 2545 (1998).
- [92] D. J. Tozer and N. C. Handy, *J. Phys. Chem. A* **102**, 3162 (1998).
- [93] N. C. Handy and D. J. Tozer, *Mol. Phys.* **94**, 707 (1998).
- [94] A. J. Thakkar and S. P. McCarthy, *J. Chem. Phys.* **131**, 134109 (2009).
- [95] L. C. Wilson, *Chem. Phys.* **181**, 337 (1994).
- [96] L. C. Wilson and S. Ivanov, *Int. J. Quantum Chem.* **69**, 523 (1998).
- [97] L. C. Wilson and M. Levy, *Phys. Rev. B* **41**, 12930 (1990).
- [98] X. Xu and W. A. Goddard, *Proc. Natl. Acad. Sci. U. S. A.* **101**, 2673 (2004).
- [99] X. Xu and W. A. Goddard, *J. Chem. Phys.* **121**, 4068 (2004).
- [100] L. A. Constantin, E. Fabiano, and F. Della Sala, *Phys. Rev. B* **84**, 233103 (2011).
- [101] L. A. Constantin, E. Fabiano, and F. D. Sala, *J. Chem. Phys.* **137**, 194105 (2012).
- [102] E. Fabiano, L. A. Constantin, P. Cortona, and F. Della Sala, *J. Chem. Theory Comput.* **11**, 122 (2015).
- [103] A. D. Becke, *J. Chem. Phys.* **88**, 1053 (1988).
- [104] A. D. Becke, *Int. J. Quantum Chem.* **52**, 625 (1994).
- [105] A. D. Becke, *J. Chem. Phys.* **104**, 1040 (1996).
- [106] A. C. Cancio and M. Y. Chou, *Phys. Rev. B* **74**, 081202 (2006).
- [107] R. Colle and O. Salvetti, *Theor. Chim. Acta* **37**, 329 (1975).
- [108] P. Verma and D. G. Truhlar, *J. Phys. Chem. C* **121**, 7144 (2017).
- [109] S. Lehtola and M. A. L. Marques, *J. Chem. Theory Comput.* **17**, 943 (2021).
- [110] J. Rey and A. Savin, *Int. J. Quantum Chem.* **69**, 581 (1998).
- [111] J. B. Krieger, J. Chen, G. J. Iafrate, and A. Savin, Construction of an accurate self-interaction-corrected correlation energy functional based on an electron gas with a gap, in *Electron Correlations and Materials Properties*, edited by A. Gonis, N. Kioussis, and M. Ciftan (Springer US, Boston, MA, 1999) pp. 463–477.
- [112] J. B. Krieger, J. Chen, and S. Kurth, *AIP Conf. Proc.* **577**, 48 (2001).
- [113] S. Kurth, J. P. Perdew, and P. Blaha, *Int. J. Quantum Chem.* **75**, 889 (1999).
- [114] J. Toulouse, A. Savin, and C. Adamo, *J. Chem. Phys.* **117**, 10465 (2002).
- [115] C. Lee and R. G. Parr, *Phys. Rev. A* **42**, 193 (1990).
- [116] Y. Zhao and D. G. Truhlar, *J. Chem. Phys.* **125**, 194101 (2006).
- [117] Y. Zhao and D. G. Truhlar, *Theor. Chem. Acc.* **120**, 215 (2008).
- [118] R. Peverati and D. G. Truhlar, *J. Phys. Chem. Lett.* **3**, 117 (2012).
- [119] R. Peverati and D. G. Truhlar, *Phys. Chem. Chem. Phys.* **14**, 13171 (2012).
- [120] H. S. Yu, X. He, and D. G. Truhlar, *J. Chem. Theory Comput.* **12**, 1280 (2016).
- [121] J. P. Perdew, S. Kurth, A. Zupan, and P. Blaha, *Phys. Rev. Lett.* **82**, 2544 (1999).
- [122] D. Mejía-Rodríguez and S. B. Trickey, *Phys. Rev. B* **102**, 121109 (2020).
- [123] J. W. Furness, A. D. Kaplan, J. Ning, J. P. Perdew, and J. Sun, *J. Phys. Chem. Lett.* **11**, 8208 (2020).
- [124] J. W. Furness, A. D. Kaplan, J. Ning, J. P. Perdew, and

- J. Sun, *J. Phys. Chem. Lett.* **11**, 9248 (2020).
- [125] Y. Wang, X. Jin, H. S. Yu, D. G. Truhlar, and X. He, *Proc. Natl. Acad. Sci. U. S. A.* **114**, 8487 (2017).
- [126] P. D. Mezei, G. I. Csonka, and M. Kállay, *J. Chem. Theory Comput.* **14**, 2469 (2018).
- [127] S. Jana, K. Sharma, and P. Samal, *J. Phys. Chem. A* **123**, 6356 (2019).
- [128] J. P. Perdew, A. Ruzsinszky, G. I. Csonka, L. A. Constantin, and J. Sun, *Phys. Rev. Lett.* **106**, 179902 (2011).
- [129] A. P. Bartók and J. R. Yates, *J. Chem. Phys.* **150**, 161101 (2019).
- [130] D. Mejía-Rodríguez and S. B. Trickey, *Phys. Rev. A* **96**, 052512 (2017).
- [131] D. Mejía-Rodríguez and S. B. Trickey, *Phys. Rev. B* **98**, 115161 (2018).
- [132] J. Tao and Y. Mo, *Phys. Rev. Lett.* **117**, 073001 (2016).
- [133] J. Tao, J. P. Perdew, V. N. Staroverov, and G. E. Scuseria, *Phys. Rev. Lett.* **91**, 146401 (2003).
- [134] J. P. Perdew, J. Tao, V. N. Staroverov, and G. E. Scuseria, *J. Chem. Phys.* **120**, 6898 (2004).
- [135] T. V. Voorhis and G. E. Scuseria, *J. Chem. Phys.* **109**, 400 (1998).
- [136] Q. Zhao, M. Levy, and R. G. Parr, *Phys. Rev. A* **47**, 918 (1993).
- [137] A. Austin, G. A. Petersson, M. J. Frisch, F. J. Dobek, G. Scalmani, and K. Throssell, *J. Chem. Theory Comput.* **8**, 4989 (2012).
- [138] C. Adamo and V. Barone, *Chem. Phys. Lett.* **274**, 242 (1997).
- [139] D. I. Bilc, R. Orlando, R. Shaltaf, G.-M. Rignanese, J. Íñiguez, and P. Ghosez, *Phys. Rev. B* **77**, 165107 (2008).
- [140] M. T. Caldeira and R. Custodio, *J. Mol. Model.* **25**, 62 (2019).
- [141] P. J. Stephens, F. J. Devlin, C. F. Chabalowski, and M. J. Frisch, *J. Phys. Chem.* **98**, 11623 (1994).
- [142] M. Reiher, O. Salomon, and B. A. Hess, *Theor. Chem. Acc.* **107**, 48 (2001).
- [143] Defined through Gaussian implementation.
- [144] A. D. Becke, *J. Chem. Phys.* **98**, 5648 (1993).
- [145] Y. Shao, M. Head-Gordon, and A. I. Krylov, *J. Chem. Phys.* **118**, 4807 (2003).
- [146] P. J. Wilson, T. J. Bradley, and D. J. Tozer, *J. Chem. Phys.* **115**, 9233 (2001).
- [147] T. W. Keal and D. J. Tozer, *J. Chem. Phys.* **123**, 121103 (2005).
- [148] A. D. Becke, *J. Chem. Phys.* **107**, 8554 (1997).
- [149] A. D. Becke, *J. Chem. Phys.* **98**, 1372 (1993).
- [150] M. Renz, K. Theilacker, C. Lambert, and M. Kaupp, *J. Am. Chem. Soc.* **131**, 16292 (2009).
- [151] M. Kaupp, M. Renz, M. Parthey, M. Stolte, F. Würthner, and C. Lambert, *Phys. Chem. Chem. Phys.* **13**, 16973 (2011).
- [152] J. Carmona-Espíndola, J. L. Gázquez, A. Vela, and S. B. Trickey, *Theor. Chem. Acc.* **135**, 120 (2016).
- [153] Z. M. Sparrow, B. G. Ernst, T. K. Quady, and R. A. DiStasio, *J. Phys. Chem. Lett.* **13**, 6896 (2022).
- [154] C. Y. Lin, M. W. George, and P. M. W. Gill, *Aust. J. Chem.* **57**, 365 (2004).
- [155] E. Fabiano, L. A. Constantin, and F. Della Sala, *Int. J. Quantum Chem.* **113**, 673 (2013).
- [156] J. K. Kang and C. B. Musgrave, *J. Chem. Phys.* **115**, 11040 (2001).
- [157] V. Tognetti, P. Cortona, and C. Adamo, *Chem. Phys. Lett.* **439**, 381 (2007).
- [158] B. J. Lynch, P. L. Fast, M. Harris, and D. G. Truhlar, *J. Phys. Chem. A* **104**, 4811 (2000).
- [159] C. Adamo and V. Barone, *J. Chem. Phys.* **108**, 664 (1998).
- [160] Y. Zhao and D. G. Truhlar, *J. Phys. Chem. A* **108**, 6908 (2004).
- [161] W.-M. Hoe, A. J. Cohen, and N. C. Handy, *Chem. Phys. Lett.* **341**, 319 (2001).
- [162] D. N. Tahchieva, D. Bakowies, R. Ramakrishnan, and O. A. von Lilienfeld, *J. Chem. Theory Comput.* **14**, 4806 (2018).
- [163] P. Cortona, *J. Chem. Phys.* **136**, 086101 (2012).
- [164] S. Grimme, J. Antony, S. Ehrlich, and H. Krieg, *J. Chem. Phys.* **132**, 154104 (2010).
- [165] Y. A. Bernard, Y. Shao, and A. I. Krylov, *J. Chem. Phys.* **136**, 204103 (2012).
- [166] C. Adamo and V. Barone, *J. Chem. Phys.* **110**, 6158 (1999).
- [167] M. Ernzerhof and G. E. Scuseria, *J. Chem. Phys.* **110**, 5029 (1999).
- [168] Y. Jin and R. J. Bartlett, *J. Chem. Phys.* **149**, 064111 (2018).
- [169] L. Lu, H. Hu, H. Hou, and B. Wang, *Comput. Theor. Chem.* **1015**, 64 (2013).
- [170] H. L. Schmider and A. D. Becke, *J. Chem. Phys.* **108**, 9624 (1998).
- [171] R. Peverati and D. G. Truhlar, *J. Chem. Phys.* **135**, 191102 (2011).
- [172] K. W. Wiitala, T. R. Hoye, and C. J. Cramer, *J. Chem. Theory Comput.* **2**, 1085 (2006).
- [173] A. D. Becke, *J. Chem. Phys.* **109**, 2092 (1998).
- [174] Y. Zhao, B. J. Lynch, and D. G. Truhlar, *J. Phys. Chem. A* **108**, 2715 (2004).
- [175] R. Neumann and N. C. Handy, *Chem. Phys. Lett.* **246**, 381 (1995).
- [176] K. Pernal, R. Podeszwa, K. Patkowski, and K. Szalewicz, *Phys. Rev. Lett.* **103**, 263201 (2009).
- [177] J. Tao, *J. Chem. Phys.* **116**, 2335 (2002).
- [178] Y. Zhao, N. E. Schultz, and D. G. Truhlar, *J. Chem. Theory Comput.* **2**, 364 (2006).
- [179] Y. Zhao, N. E. Schultz, and D. G. Truhlar, *J. Chem. Phys.* **123**, 161103 (2005).
- [180] Y. Zhao and D. G. Truhlar, *J. Phys. Chem. A* **110**, 13126 (2006).
- [181] Y. Zhao and D. G. Truhlar, *J. Chem. Theory Comput.* **4**, 1849 (2008).
- [182] H. S. Yu, X. He, S. L. Li, and D. G. Truhlar, *Chem. Sci.* **7**, 5032 (2016).
- [183] Y. Zhao, N. González-García, and D. G. Truhlar, *J. Phys. Chem. A* **109**, 2012 (2005).
- [184] Y. Zhao and D. G. Truhlar, *J. Chem. Theory Comput.* **1**, 415 (2005).
- [185] Y. Zhao and D. G. Truhlar, *J. Phys. Chem. A* **109**, 5656 (2005).
- [186] Y. Wang, P. Verma, X. Jin, D. G. Truhlar, and X. He, *Proc. Natl. Acad. Sci. U. S. A.* **115**, 10257 (2018).
- [187] G. I. Csonka, J. P. Perdew, and A. Ruzsinszky, *J. Chem. Theory Comput.* **6**, 3688 (2010).
- [188] S. Grimme, *J. Phys. Chem. A* **109**, 3067 (2005).
- [189] Y. Zhao, B. J. Lynch, and D. G. Truhlar, *Phys. Chem. Chem. Phys.* **7**, 43 (2005).
- [190] V. N. Staroverov, G. E. Scuseria, J. Tao, and J. P. Perdew, *J. Chem. Phys.* **119**, 12129 (2003).